

Age Assignments of Larger Foraminiferal Assemblages of Maastrichtian to Eocene Age in Northern Pakistan

By

WOLFGANG WEISS^{*)}

With 12 Text-figures and 9 Plates

ABSTRACT

Geological, microfaunal and biostratigraphical records were obtained from selected sections of the Upper Indus Basin of northern Pakistan (the western Salt Range, the Kohat area, the Sulaiman Range).

Eight assemblages of larger foraminifers were identified within the Late Cretaceous and Early Paleogene formations. They were dated as exactly as possible by means of Tethyan

planktonic foraminifers. Several refinements of previous age assessments were obtained. The larger foraminifers which were identified are important for establishing a firm age of the shallow marine formations. They support the knowledge of the paleobiogeography of the larger foraminifers within the eastern Tethys realm.

ZUSAMMENFASSUNG

Es werden geologische, mikrofaunistische und biostratigraphische Ergebnisse dargestellt, die aus Untersuchungen an Referenzprofilen des oberen Indus-Becken im nördlichen Pakistan (der westlichen Salt Range, der Kohat-Region und der Sulaiman Range) gewonnen wurden.

Acht Großforaminiferen-Gesellschaften aus der jüngsten Oberkreide und dem Paläogen wurden festgestellt. Sie wurden mit Hilfe der tethyalen Standardgliederung nach planktoni-

schen Foraminiferen so genau wie möglich datiert. Dadurch können frühere Altersangaben verfeinert werden. Die Kenntnis der verschiedenen Großforaminiferen-Gesellschaften hat wesentliche Bedeutung für die Biostratigraphie der flachmarinen Sedimente der Region. Die Beobachtungen bekräftigen die Kenntnisse zur paläobiographischen Verbreitung der Großforaminiferen der östlichen Tethys.

1. INTRODUCTION

The author participated in several reconnaissance field parties in the Upper Indus Basin of northern Pakistan in 1985 and 1986. The biostratigraphy and paleoenvironment of late Mesozoic and Paleogene formations of the Upper Indus Basin, in particular that of the Salt Range, the Kohat area and the Sulaiman Range, were studied on the basis of microfossils, with preference on planktonic and larger foraminifers (WEISS 1988).

The main objectives of the microfaunal and foraminiferal studies were (1) to compare Cretaceous and Paleogene reference sections in several areas of northern Pakistan (the western Salt Range, the Kohat area and the Sulaiman Range), (2) to ascertain the biostratigraphic ages of the Mesozoic and Cenozoic formations and express these ages preferably in terms of modern standard planktonic foraminiferal zonation, and (3) to describe the widely distributed assemblages of larger foraminifers of the different Upper Indus formations and to integrate these assemblages as exactly as possible into the standard Cretaceous and Cenozoic planktonic foraminiferal zones.

^{*)} Dr. WOLFGANG WEISS, Bundesanstalt für Geowissenschaften und Rohstoffe (BGR), Stilleweg 2, D-30655 Hannover




Fig. 1: Location map of investigated sections of the Upper Indus Basin (after EAMES 1952, Geological Map of Pakistan 1:2 000 000 1964, SIIAH 1977 and PORTH & KAMRAN in PORTH & HILAL A. RAZA 1988).

The distribution of the larger foraminifers of the Upper Cretaceous and Paleogene presented in this paper is based on regional standard sections (Fig. 1) which were more or less continuously sampled. Samples would be taken preferably at the base and the top of the formations and at several horizons within the formations which were found to reflect typical lithologies or were found remarkable either through changing lithology or through different faunal contents as seen from visual inspections.

The studied intervals are from the following sections and areas:

- the western Salt Range:

- (1) the Paleogene interval of the Patala Nala section;
- (2) the Paleogene interval of the Nammal Gorge section;
- (3) the Paleogene interval of the Khairabad section;

the Kohat area:

- (4) the Cretaceous to Paleogene interval of the section North of Hangu;
- (5) the Paleogene interval of the Panoba Anticline section;

- the Sulaiman Range:

- (6) the Jurassic to Paleogene interval of the Mughal Kot-Toi Nala section;
- (7) the Paleogene interval of the Zinda Pir section;
- (8) the Eocene interval of the Zao River section;
- (9) the Cretaceous to Paleogene interval of the Rakhi Nala section;
- (10) the Cretaceous to Paleogene interval of the Murrey Brewery Gorge section.

About 158 microfaunas from washed residues of shales and marls and about 100 thin sections from limestone samples are

reported here. Stratigraphic assignments and formation determinations of the studied intervals of the sections were done mainly according to the literature (SHAH 1977, PORTH & HILAL A. RAZA 1990a, b) and field observations. Composition of microfaunas and in particular the distribution of planktonic and larger foraminifers which were investigated in detail are given in Figs. 2-11.

Cretaceous and Paleogene foraminiferal taxonomy and zones are adopted mainly from BOLLI (1957, 1966), BLOW (1969, 1979), POSTUMA (1971), STAINFORTH et al. (1975), ROBASZYNSKI & CARON (1979), ROBASZYNSKI et al. (1984), CARON (1985), and TOUMARKINE & LUTERBACHER (1985) for the planktonic foraminifers, and from DAVIES & PINFOLD (1937), SMOOTH (1954), NAGAPPA (1959), HOTTINGER (1960), ADAMS (1970), KURESHI (1975, 1978) and SCHAUB (1981) for the larger foraminifers.

ACKNOWLEDGEMENTS

I wish to thank the staff of the Hydrocarbon Development Institute of Pakistan (HDIP) (Mr. HILAL A. RAZA, Director General and Head of the HDIP, Mr. RIAZ AHMED, Principal Geologist of Field Geology and Petroleum Geology, Mr. MUZAFFAR KAMRAN, Senior Geologist, Mr. IDREES KHAN, Senior Chemist, Mr. SALIM MEHMOOD, Geologist) and the project leaders of the Federal Institute for Geosciences and Natural Resources (BGR) (Dr. PORTH, Dr. RANKE, Dr. JURGAN) who shared the days in the field with me and for stimulating

discussions. I also thank the people of the Pakistan provinces and tribes for their hospitality. I am grateful to Mr. LUPPOLD and Mrs. ENGEL (BGR) for sample preparation and drafting. Dr. WOLFART (BGR) is thanked for his critical reading an earlier version of the manuscript. My colleagues Mrs. Dr. FENNER, Mrs. Dr. KOTHE, Dr. VON DANIELS and Dr. GRAMANN have offered constructive criticism of the manuscript and many important suggestions.

2. GEOLOGICAL SETTING

The Indus Basin of Pakistan is subdivided in two basins, the Upper Indus Basin and the Lower Indus Basin, each consisting of several subbasins, ranges, plateaus, provinces (SHAH 1977, PORTH & HILAL A. RAZA 1990a, b). The various structural elements which are arranged in a garland-like pattern along the Indo-Pakistan-Eurasian plate convergence in Pakistan belong to different tectonic units, the Hazara Block and the Sulaiman Block, of the Indo-Pakistan Plate which is separated by the Makran-Khojak-Pishin Flysch Belt from the Afghanistan and Turan Blocks of the Eurasian Plate (BANNERT 1992: fig. 1).

The locations of the studied sections of the Upper Indus Basin (the western Salt Range, the Kohat area, the Sulaiman Range) which belong to the Hazara Block and the Sulaiman Block of the Indo-Pakistan Plate are shown in Fig. 1.

Mesozoic and Paleogene sediments of the eastern fold belts of the Upper Indus Basin (as well as of the Lower Indus Basin) are deposited on a broad shelf area of the passive continental margin of the Indo-Pakistan Plate (BANNERT 1992). The fold belts originated since the Paleocene as a result of the main

collision of the Indo-Pakistan Plate with the Eurasian Plate (BANNERT 1992: 7).

The formations of the Salt Range comprise successions of stable-platform, marine sediments of Late Precambrian and Cambrian, Permian, Triassic, Jurassic, Cretaceous (questionable), Paleogene age, followed by Early Miocene to Pleistocene molasse sediments (GEE 1980, 1989).

In the northwestern Kohat area (the Kohat Salt Region) the surface formations are equally of shallow marine platform type and are of Tertiary to Quaternary age. Late Cretaceous sediments with poor ammonite and foraminiferal biostratigraphy (LATIF 1970, FATMI 1973, SHAH 1977) are exposed occasionally in the cores of the anticlines.

In the Sulaiman Range shallow marine shelf sediments have been documented for the Permian to the Late Eocene (SHAH 1977). Paleogene shallow marine sediments of the Sulaiman Range show some differences in both facies and biostratigraphy from those of the western Salt Range and the Kohat area, reflected by lithology or microfauna.

3. RESULTS

3.1 REFINEMENTS OF THE REGIONAL STRATIGRAPHY

The succession and microfaunal biostratigraphy of the Mesozoic and Cenozoic formations of the Upper Indus Basin of northern Pakistan has been summarized by a number of authors which gave a large amount of informations (e. g. NUTTALL 1925, 1926, DAVIES 1926, 1927, DAVIES & PINFOLD 1937, EAMES 1952, GILL 1952, 1953, HAQUE 1956, SMOUT & HAQUE 1956, NAGAPPA 1959, HUNTING SURVEY CORPORATION 1961, MARKS 1962, LATIF 1963, 1964, ADAMS 1970, FATMI 1973, KURESHY 1966, 1969, 1971, 1972, 1974, 1975, 1976, 1977, 1978, SHAH 1977, PORTH & HILAL A. RAZA 1990a, b). New biostratigraphic results are available from studies on calcareous nannofossils and dinoflagellates by KOTHE (1987, 1988), of benthic and planktonic foraminifers by BUTT (1987, 1991), NASEER A. SHAFIQUE & VON DANIELS (1990) and JAMIL AFZAL & VON DANIELS (1991).

The revised nomenclature for the stratigraphic units of the Cretaceous and Paleogene in northern Pakistan, set up by the Stratigraphic Committee of Pakistan (FATMI 1973, SHAH 1977), is followed here with minor modifications.

Biostratigraphic age assignments of the different Upper Cretaceous and Paleogene formations and related lithostratigraphic units on the basis of foraminiferal data from the author's studies are shown in Figs. 2-11.

The following results should be mentioned:

1) According to planktonic foraminifers, the beginning of the pelagic Goru Formation in the Mughal Kot-Toi Nala section and the Murree Brewery Gorge section of the Sulaiman Range (Figs. 7-8) was dated as late Early Cretaceous and equivalent to the planktonic foraminiferal *Rotalipora tucinensis* and *Rotalipora appenninica* Zones of Late Albian. According to planktonic foraminifers described by GIGON (1962) and




Fig. 2: Distribution of larger and planktonic foraminifers of the Patala Nala section, western Salt Range.

Abbreviations used in Figs. 2-12.

- Microfauna: rad = radiolarians
 pF2 = planktonic foraminifers
 bF = benthic foraminifers
 lF = larger foraminifers
 os = ostracods
 al = algae
 mf = mesofauna

Planktonic foraminiferal genera: A. = *Acarinina*, Ab. = *Abatbomphalus*, „G.“ = *Globigerina*, G. = *Globotruncana*, Gt. = *Globigermatheka*, H. = *Hantkenina*, M. = *Morozovella*, O. = *Orbulinoides*, Pl. = *Planorotalites*, T. = *Turborotalia*, Tr. = *Truncorotaloides*.

Larger foraminiferal genera: Ass. = *Assilina*, D. = *Discocyclina*, Dictyoc. = *Dictyocenoidea*, Dictyoc. = *Dictyocatabina*, Fl. = *Flosculina*, L. = *Lockbartia*, Misc. = *Miscellanea*, N. = *Nummulites*, Omphaloc. = *Omphalocyclus*, Orb. = *Orbitoides*, Rot. = *Rotalia*, Sid. = *Siderolites*.

Cretaceous planktonic foraminiferal zones according to CARON (1985), Paleogene planktonic foraminiferal zones according to TOUMARKINE & LUTERBACHER (1985).

Lithological symbols used in Figs. 2-11

- = conglomerate
- = sandstone, partly calcareous
- = claystone, shale, partly sandy or calcareous
- = limestone
- = marl
- = gypsum




Fig. 3: Distribution of larger and planktonic foraminifers of the Namgal Gorge section, western Salt Range. Abbreviations see Fig. 2.




Fig. 4: Distribution of larger and planktonic foraminifers of the Khairabad section, western Salt Range. Abbreviations see Fig. 2.

FRITZ & KHAN (1967), the upper limit of the Goru Formation could be assigned within an interval between the Early-Middle Cenomanian *Rotalipora reicheli* Zone to Early/Middle Campanian *Globotruncana ventricosa* Zone of planktonic foraminifers.

There are no supporting data by the author's studies based on planktonic foraminifers proving the correlation of the Goru Formation (Late Albian to Late Turonian-Early Coniacian) of the Sulaiman Range with the generally unfossiliferous Lumshival Sandstone Formation of the western Kohat area (SHAH 1977: 54) which was assigned to be of Aptian-Early Albian(!) age. The Lumshival Sandstone Formation underlies the Late Cretaceous Darsamad Limestone of the Kawagarh Formation (PORTH & HILAL A. RAZA 1990a: fig. 4; PORTH & HILAL A. RAZA 1990b: fig. 2) (see below).

2) There is a remarkable facies and paleoenvironment change between the Parh Formation which is lithologically similar to the underlying Goru Formation (HUNTING SURVEY CORPORATION (1961) and KURESHI (1978) included these strata in the Parh Group!) and the overlying Mughal Kot Formation, as seen in the Mughal Kot-Toi Nala section (Fig. 7; compare also EAMES 1952, WILLIAMS 1959, SHAH 1977). The change of facies

was dated, according to planktonic foraminifers, as close to Late Turonian to Early Coniacian times (see below).

The Parh Formation consists of pelagic, light-grey limestone and marlstone sequences (about 300 m thick) and is similar to that of the Goru Formation (about 120 m thick), whereas the Mughal Kot Formation (about 1150 m thick; thickness according to SHAH 1977 and PORTH 1990b) comprises conglomerates, thick shale and marlstone sequences. Several sequences are showing slumping and gliding, as repeatedly observed during field parties in the Mughal Kot-Toi Nala and the Rakhi Nala sections. The facies change from pelagic ocean floor sediments to flyschoid foredeep sediments is indicated by microfossils by the complete lack of autochthonous foraminifers within lower parts of the Mughal Kot Formation. No foraminiferal data were obtained from this formation except for scarce reworked and fragmented foraminiferal tests. Most of the samples are completely barren of foraminifers (compare Fig. 7).

From the Parh Formation one sample (SmR-131) taken 5 m below the top of the formation yielded planktonic foraminifers with *Margino-truncana pseudolimneciana* PESSAGNO and *Heterobelix* specimens close to *Heterobelix morenami* (CUSHMAN)




Fig. 5: Distribution of larger and planktonic foraminifers of the North of Hangu section, Kohat area. Abbreviations see Fig. 2.




Fig. 6: Distribution of larger and planktonic foraminifers of the Panaba Anticline section, Kohat area. Abbreviations see Fig. 2.

and *Heterohelix reussi* (CUSHMAN). Another sample (SmR-132) from about 30 m below the top has shown poor planktonic foraminifers from which specimens of the high-spired *Praeglobotruncana turbinata* (REICHEL) were identified. The last occurrence of high-spired praeglobotruncanids is within the Middle Turonian, whereas the range of *Marginitruncana pseudolinneiana* PESSAGNO is from Middle Turonian to Early Coniacian according to ROBASYNSKI & CARON (1979).

Therefore, the top of the Park Formation of the Mughal Kot-Toi Nala section of the Sulaiman Range was dated as belonging to the planktonic foraminiferal *Marginitruncana sigali* to *Marginitruncana primitiva* Zones (Fig. 7) which are of Late Turonian to Early Coniacian according to CARON (1985: fig. 3).

3) In the Murree Brewery Gorge section of the Sulaiman Range (Fig. 11), the limestones of the pelagic Goru and Park




Fig. 7. Distribution of larger and planktonic foraminifers of the Mughal Kot-Toi Nala section, Sulaiman Range. Abbreviations see Fig. 2.




Fig. 8. Distribution of larger and planktonic foraminifers of the Zinda Pir section, Sulaiman Range. Abbreviations see Fig. 2.

Formations show low sedimentation rates (the limestones are altogether about 120 m thick) and contain well-preserved and abundant planktonic foraminifers, such as globotruncanids, hedbergellids and heterohelicids. Planktonic foraminiferal zones from the Late Albian *Rotalipora appenninica* Zone to the Late Santonian *Marginotruncana asymetrica* Zone have been identified (Fig. 11). They confirm former age determinations by ALLEMANN (1979). The transition from the Goru Formation to the Parh Formation is either within the interval of the *Marginotruncana sigali* and the *Marginotruncana primitiva* Zones or within the interval of the *Marginotruncana primitiva* and *Marginotruncana concavata* Zones. The age assignment is

Early to Late Coniacian according to CARON (1985: fig. 3). The transition is characterized by the disappearance of high trochospiral praeglobotruncanids, i. e. *Praeglobotruncana turbinata* (REICHEL), and by the appearance of early globotruncanids, i. e. *Globotruncana fomicata* PLUMMER.

All limestones from the Murree Brewery Gorge section contain highly diverse and abundant planktonic foraminifers with many keeled species, the sequence must have originated in a tropical oceanic facies. The pelagic part of the Upper Cretaceous sequence of the Murree Brewery Gorge section represents „the floor of Neo-Tethyan ocean between the northern margin of the Indo-Pakistan Plate and the southern




Fig. 11: Distribution of larger and planktonic foraminifers of the Murree Brewery Gorge section, Sulaiman Range. Abbreviations see Fig. 2.

intervals. In the Mughal Kot-Toi Nala section and the Rakhi Nala section of the Sulaiman Range (Figs. 7, 10), the Paleocene-Eocene boundary expected within the Ghazij Shales and between the Lower and Upper Rakhi Gaj Shales (SHAH 1977: 70) was not definitely recognizable due to very bad preservation of morozovellids.

8) The Early Eocene part of the Brewery Limestone in the Murree Brewery Gorge section of the Sulaiman Range (Fig. 11) is characterized by a *Nummulites globulus-Operculina salsa* Assemblage of larger foraminifers. According to BLONDEAU (1972) and SCHAUB (1981), *Nummulites globulus* LEYMERIE is restricted to Early Eocene, whereas according to BAYLIS (1961: 97/98) *Nummulites globulus* LEYMERIE is present up to the Middle Eocene of the Kirthar Formation.

The Early Eocene part of the Brewery Limestone (Fig. 11) correlates (a) with the upper part of the Zinda Pir Limestone which shows a characteristic *Alveolina-Nummulites fossulata* Assemblage (Fig. 8), and (b) with parts of the Gazij Formation which are characterized by the *Orbitolites complanatus-Discocyclina ranikotensis-Nummulites fossulata* Assemblage (Fig. 7). These assemblages of larger foraminifers which were not found outside of the Sulaiman Range contain the first small, but definitive *Nummulites* specimens with *Nummulites globulus* LEYMERIE as the most common species and are assigned to be of Early Eocene (see above).

9) The Green Shales and the Nodular Shales of EAMES (1952) from the Rakhi Nala section of the Sulaiman Range (Fig. 10) are characterized by an *Assilina leymeriei-Nummulites fossulata* Assemblage of larger foraminifers. *Nummulites fossulata* DE CIZANCOURT is known from Early Eocene according to this author and was found to be a larger foraminiferal index marker for the Early Eocene interval of the Rakhi Nala section. Based on the characteristic larger foraminiferal assemblage, the Green and Nodular Shales could be correlated with basal parts of the Kirthar Formation of the Mughal Kot-Toi Nala section (Fig. 7) which contains an *Assilina spinosa-Nummulites ataicus* Assemblage as well as with the „*Assilina cliff*“ of the Zao River section (Fig. 9). These assemblages of larger foraminifers are further characterized by specimens of *Discocyclina ranikotensis* DAVIES which disappears at the top of the assemblage (Fig. 12).

10) The Early-Middle Eocene boundary was expected at the base of Platy Limestone of the Rakhi Nala section (Fig. 10; and

EAMES 1952) or between the less prominent Early Eocene „*Assilina cliff*“ and the prominent Middle Eocene „*Discocyclina cliff*“ of the Mughal Kot-Toi Nala and Zao River sections of the Sulaiman Range (Figs. 7, 9). The „*Discocyclina cliff*“ which might be paleoenvironmentally a submarine barrier is named after characteristic field morphology and composition of limestones consisting of abundant, sometimes monospecific *Discocyclina* specimens (*Discocyclina dispansa* (SOWERBY), *Discocyclina sowerbyi* NUTTALL, and - less common - *Discocyclina undulata* NUTTALL and *Discocyclina dorreini* BAYLIS). The *Discocyclina* specimens are associated often with *Alveolina elliptica* (SOWERBY) and *Planorbulina* sp. The „*Discocyclina cliff*“ was dated with planktonic foraminifers and is within the interval from the *Globigerinatheka subconglobata* to the *Orbulinoides beckmanni* Zones according to TOUTMARKINE & LUTERBACHER (1985: fig. 1).

3.2 ASSEMBLAGES OF LARGER FORAMINIFERS

Because of paleoenvironmental and paleobiogeographical limitations, there is no standard zonation accepted world-wide for the larger foraminifers of the Mesozoic and Cenozoic. There are several regional and interregional successions of Cretaceous and Tertiary larger foraminifers. The most relevant to the Indus Basin are the Late Cretaceous orbitoidal foraminiferal zonation (VAN GORSEL 1978), the Tertiary Letter Classification (ADAMS 1970), the Tertiary larger foraminiferal zones of Pakistan (KURESHY 1978) and the phylogeny of various lineages of Paleogene alveolinids and operculinids, nummulitids and assilininids, as described by HOTTINGER (1960, 1977), BUTT (1987, 1991), BLONDEAU (1972) and SCHAUB (1981). They include detailed microfaunal and stratigraphical informations of earlier workers (e. g. NUTTALL 1925, 1926; DAVIES 1937; DE CIZANCOURT 1938; GILL 1952, 1953; SMOOTH 1954; SMOOTH & HAQUE 1956; NAGAPPA 1959; MARKS 1962; SEN GUPTA 1963).

The larger foraminiferal associations which were identified during this study in the Upper Cretaceous and Paleogene sections of northern Pakistan are understood here as assemblages of paleoenvironmental significance (compare NICORA et al. 1986).

The Paleogene assemblages may be assigned to „Tertiary a1 (Late Paleocene)“, the „Tertiary a2 (Early Eocene)“ and

„Tertiary a3 (Middle Eocene)“ of the Letter Stage Classification (ADAMS 1970). Correlation to the „Lower Paleocene (Danian) *Daviesina langhami* Zone“, the „Upper Paleocene (Landenian) *Nummulites nuttalli* Zone“, the „Lower Eocene (Ypresian) *Assilina granulosa* Zone“ and the „Middle Eocene (Lutetian) *Nummulites beaumonti* Zone“ of KURESHY (1978) is possible, but less precise due to differences in species assemblages.

The biostratigraphic assignment of larger foraminiferal assemblages was determined following HAAK & POSTUMA (1975), CARON (1985) and TOUMARKINE & LUTERBACHER (1985) on the basis of the associated planktonic foraminifers with which larger foraminifers were found in some favourable instances. The co-occurrence of larger and planktonic foraminifers indicates in general a paleoenvironment within outer parts of the Sublittoral. It was not possible in any case to assign individual larger foraminifer species precisely to planktonic foraminiferal zones. The natural habitat and deposition area of most of the larger foraminiferal specimens are shallow-water paleoenvironments which are not preferred by planktonic foraminifers. These paleoenvironmental differences are the main reason for uncertainties in the positions of the first and last occurrences of larger foraminiferal species of Fig. 12.

However, it was found by several authors (cit. below) that some species of larger foraminifers, such as *Omphalocyclus macroporus* (LAMARCK), *Siderolites calcitrapoides* LAMARCK, *Assilina dandotica* DAVIES, *Discocyclina ranikotensis* DAVIES, *Discocyclina dispansa* (SOWERBY) and *Discocyclina sowerbyi* NUTTALL, have a restricted distribution both in facies and biostratigraphy. These species are of importance for the determination of the biostratigraphical age of assemblages containing larger foraminifers only.

Several sections revisited and sampled were investigated earlier by DAVIES & PINFOLD (1937) (the Nammal Gorge section), EAMES (1952) (the Rakhi Nala section), BAYLISS (1961) (the Rakhi Nala section), MARKS (1962) (the Rakhi Nala section), SAMANTA (1972) (the Rakhi Nala section), LATIF (1972) (the section North of Hangu), ALLEMAN (1979) (the Murree Brewery Gorge section) and (GEE 1980) (other sections of the Salt Range and the Kohat area).

The distribution of larger foraminifers of Late Cretaceous and Early Paleogene as derived from the above mentioned regional sections is shown in Fig. 12.

The following assemblages of larger foraminifers have been identified and are described as follows.

3.2.1 Cretaceous

Orbitoides media-Omphalocyclus macroporus Assemblage

Age: Early Maastrichtian

Remarks: This assemblage of larger foraminifers is characterized by the co-occurrence of *Orbitoides media* (D'ARCHIAC), *Orbitoides apiculata* SCHLUMBERGER (which represents the final stage of the *Orbitoides* lineage (VAN GORSEL 1978: 31), *Orbitoides tissoti* SCHLUMBERGER, *Omphalocyclus macroporus* (LAMARCK) and *Siderolites calcitrapoides* LAMARCK. Variations of subspecies of *Orbitoides tissoti* SCHLUMBERGER

from the Late Cretaceous of the Rakhi Nala section are described in detail by MARKS (1962).

The assemblage was found typically within the *Orbitoides* Limestone sequence of the Rakhi Nala section of the Sulaiman Range (Fig. 10). It is consistent with the „*Orbitoides* limestones and shales“ of EAMES (1952) which were subdivided later (SHAH 1977) in a lower part assigned to the Mughal Kot Formation and an upper part assigned to the Fort Munro Formation. Due to the evident conformity based on larger foraminiferal assemblage, this part of the Fort Munro Formation which represents the upper part of „*Orbitoides* limestones and shales“ of EAMES (1952) is re-included in the Mughal Kot Formation.

The *Orbitoides media-Omphalocyclus macroporus* Assemblage is dominated by abundant *Orbitoides* specimens. It indicates a shallow water environment and may be assigned to middle parts of the Sublittoral (MARKS 1962). *Omphalocyclus macroporus* (LAMARCK) is probably a marker sensitive to successive shallowing of sea level; it was found throughout the *Orbitoides* Limestone sequence but in less abundance within upper parts of the *Orbitoides* Limestone sequence underlying the near-shore, unfossiliferous Pab Sandstone Formation.

The stratigraphic distribution of the above mentioned *Orbitoides* species and in particular that of *Omphalocyclus macroporus* (LAMARCK) seems to be highly restricted and may be assigned approximately to the Early Maastrichtian (VAN GORSEL 1978).

Siderolites calcitrapoides-Orbitoides media Assemblage

Age: early Late Maastrichtian

Remarks: This assemblage is characterized by abundant specimens of *Siderolites calcitrapoides* LAMARCK co-occurring with thickly lenticular specimens of *Orbitoides media* (D'ARCHIAC), *Orbitoides tissoti* SCHLUMBERGER and *Orbitoides apiculata* SCHLUMBERGER. *Omphalocyclus macroporus* is not present probably due to paleoenvironmental reasons (see above). The gradual increase of the mean relative thickness of *Orbitoides* specimens is probably a response to increased water energy (compare MARKS 1962).

The assemblage was recognized in the basal Upper Cretaceous part of the Brewery Limestone within the Murree Brewery Gorge section of the Sulaiman Range (Fig. 11). It indicates a shallow water environment of middle to inner parts of the Sublittoral below or near the wave base.


The above mentioned larger foraminiferal association is known from the European Late Maastrichtian (CONCLUSIONS/NICEF 1983: 420, tab. 5b).

Siderolites calcitrapoides-Rotalia cf. trochidiformis Assemblage

Age: late Late Maastrichtian

Remarks: This larger foraminiferal association was found at the top of Upper Cretaceous sediments and contains rare specimens of *Siderolites calcitrapoides* LAMARCK and abundant specimens of *Rotalia cf. trochidiformis* (LAMARCK). Similar observations are known from the Qatar Peninsula (SMOUTH 1954).

The assemblage was observed in the topmost Upper Cretaceous part of the Brewery Limestone within the Murree



WEISS, W.: Maastrichtian to Eocene foraminiferal assemblages of Pakistan

Plate 1

Plate 1 Nammal Gorge section, Salt Range. Larger foraminifers.

- Fig. 1 Algal-larger foraminiferal assemblage with *Operculina salsa* DAVIES, *Miscellanea miscella* (D'ARCHIAC & HAIME) and others. - SaR-39/S 1317; about 20 x; Middle Paleocene.
- Fig. 2 *Lockbartia haimei* (DAVIES) and questionable *Cibicides* specimens within a sandy matrix. - SaR-40/S 1473; about 64 x; Middle Paleocene.
- Fig. 3 *Discocyclina vanikotensis* DAVIES (juvenile specimen, probably A generation) and algal fragments. - SaR-39/S 1317; about 40 x; Middle Paleocene.
- Fig. 4 *Lockbartia* aff. *conditi* (NUTTALL). - SaR-48/S 1323; about 48 x; Early to Middle Paleocene.
- Fig. 5 *Dictyokathina simplex* SMOOTH. Note the thickened dorsal side and the layered ventral side. - SaR-42/S 1319; about 48 x; Early to Middle Paleocene.
- Fig. 6 *Miscellanea miscella* (D'ARCHIAC & HAIME). - SaR-45/S 1321; about 48 x; Early to Middle Paleocene.

Brewery Gorge section of the Sulaiman Range (Fig. 11) and in uppermost parts of the massive Darsamand Limestone within the section North of Hangu in the Kohat area (Fig. 5). It indicates a very shallow paleoenvironment of inner parts of the Sublittoral.

In the section North of Hangu the assemblage is well above the base of the Late Maastrichtian *Abathomphalus mayaroensis* Zone of planktonic foraminifers and, therefore, may be assigned to the late Late Maastrichtian.

3.2.2 Paleocene

Miscellanea-Lockhartia haimiei-Dictyokathina simplex Assemblage

Age: late Early to Middle Paleocene

Remarks: This assemblage of larger foraminifers is characteristic of the Lockhart (= Khairabad) Limestone sequence which contains a rich benthic foraminiferal assemblage (DAVIES & PINFOLD 1937, HAQUE 1956). It is characterized mainly by abundant *Miscellanea miscella* (D'ARCHIAC & HAIME) and common *Lockhartia* specimens close to *Lockhartia haimiei* (DAVIES). The assemblage is additionally characterized by species which occur sporadically, such as *Kathina major* SMOUTH, *Kathina selveri* SMOUTH, *Dictyokathina simplex* SMOUTH and *Discocyclina ranikotensis* DAVIES. The occurrence of these species may reflect minor regional differences of paleoenvironmental conditions or depositional depths of the Lockhart (=Khairabad) Limestone Formation. *Dictyokathina simplex* SMOUTH seems to be restricted to this interval.

The assemblage is accompanied by *Operculina salsa* DAVIES, the probable ancestor of *Ranikothalia* species (HOTTINGER 1977, BUTT 1991). *Discocyclina ranikotensis* DAVIES is present as early as the Early-Middle Paleocene transition, if the transition coincides with the facies change from the Dhak Pass Beds into the Lockhart Limestone Formation of the Nammal Gorge section. *Orbitolites complanatus* LAMARCK reaches its first abundance maximum during this interval.

The assemblage was observed with modifications (mentioned above) in the North of Hangu section of the Kohat area (Fig. 5) and the Nammal Gorge and Khairabad sections of the western Salt Range (Figs. 3-4). It was found in uppermost parts of the Dhak Bass Beds of the Nammal Gorge section (Fig. 3).

The *Miscellanea-Lockhartia haimiei-Dictyokathina simplex* Assemblage has not been related to standard planktonic foraminiferal zonation directly due to the lack of planktonic index species. SHAH (1977) places the Lockhart Limestone Formation within the Paleocene. The base of the overlying Patala Shales at the Khairabad section (Fig. 4) could be dated as belonging to the *Planorotalites pseudomenardii* Zone of the basal Late Paleocene. The underlying Hangu Formation (=Dhak Bass Beds) was dated as Early Paleocene (SHAH 1977). Therefore, this assemblage of larger foraminifers which is typical of the Lockhart (=Khairabad) Limestone sequence may be assigned approximately to the Middle Paleocene or to an interval ranging from late Early Paleocene to Middle Paleocene, if uppermost parts of the Dhak Pass Beds are included.

Assilina dandotica-Discocyclina ranikotensis Assemblage

Age: Late Paleocene

Remarks: The assemblage is well defined biostratigraphically on the basis of planktonic foraminiferal *Planorotalites pseudomenardii* and *Morozovella velascoensis* Zones of the Late Paleocene (Figs. 2-3). It is characterized by the first appearance and abundance of true *Assilina* specimens (*Assilina dandotica* DAVIES, *Assilina leymeriei* D'ARCHIAC & HAIME, *Assilina spinosa* DAVIES) and by a first abundance maximum of *Discocyclina ranikotensis* DAVIES. It is associated with *Operculina salsa* DAVIES, *Operculina patalensis* DAVIES, not specified *Ranikothalia* specimens (see faunal reference list below) and rare specimens of *Lockhartia haimiei* (DAVIES), *Kathina major* SMOUTH and *Kathina selveri* SMOUTH. The assemblage is accompanied by small benthic and planktonic foraminifers. It indicates a deeper water environment and may be assigned to outer parts of the Sublittoral.

Assilina dandotica DAVIES seems to be a larger foraminiferal marker species for the Late Paleocene Patala Shales of the Salt Range (DAVIES & PINFOLD 1937) or the upper „Ranikot Group“ (SCHAUB 1981: 206) as defined by SHAH (1977).

The best occurrence of this larger foraminiferal assemblage was observed in the Patala Shales of the Patala Nala and the Nammal Gorge sections of the western Salt Range (Figs. 2-3) with the exception of the Khairabad section (Fig. 4) where the Patala Shales contain high percentages of planktonic foraminifers (JAMIL AFZAL & VON DANIELS 1991). In the Patala Shales from the Patala Nala type locality (Fig. 2) the assemblage of larger foraminifers occurs in higher parts of the Patala Shales, whereas lower parts of the Patala Shales are characterized by a relatively rich assemblage of small benthic foraminifers with dominant *Cibicides* and *Elphidium* specimens (called *Cibicides-Elphidium* Assemblage).

In the *Assilina dandotica-Discocyclina ranikotensis* Assemblage the *Miscellanea-Ranikothalia-Discocyclina ranikotensis* Assemblage from the lower part of the Zinda Pir Limestone (Fig. 8) is included due to planktonic foraminifers of the Late Paleocene *Morozovella velascoensis* Zone.


The *Lockhartia haimiei-Discocyclina ranikotensis* Assemblage which was identified in basal parts of the Paleogene Brewery Limestone of the Murrey Brewery Gorge section (Fig. 11) represents, according to HOTTINGER (1978) (cit. by ALLEMAN 1979: 217), the *Alveolina cucumiformis* Zone of the base of the Late Paleocene which correlates now, according to SCHAUB (1981), with the Late Paleocene *Morozovella velascoensis* Zone of planktonic foraminifers. The *Lockhartia haimiei-Discocyclina ranikotensis* Assemblage is included in the *Assilina dandotica-Discocyclina ranikotensis* Assemblage.

3.2.3 Eocene

Assilina leymeriei-Nummulites fossulata-Discocyclina ranikotensis Assemblage

Age: Early Eocene

Remarks: This assemblage is a composite of several time-equivalent associations of larger foraminifers from the studied Upper Indus sections and includes



WEISS, W.: Maastrichtian to Eocene foraminiferal assemblages of Pakistan

Plate 2

Plate 2 Nammal Gorge section, Salt Range. Larger foraminifers, Early Eocene.

- Fig. 1 Larger foraminiferal assemblage with *Discocyclina ranikotensis* DAVIES, *Discocyclina* sp., *Assilina laminosa* GILL, an intermediate specimen between *Assilina leymeriei* D'ARCHIAC & HAIME and *Assilina spinosa* DAVIES, *Nummulites globulus* LEYMERIE, *Lockbartia haimeii* (DAVIES) and others. - SaR-32/S 1315; about 12 x.
- Fig. 2 *Discocyclina ranikotensis* DAVIES, *Discocyclina* sp. and others. Note the different thicknesses at the poles. - SaR-32/S 1315; about 30 x.
- Fig. 3 *Nummulites globulus* LEYMERIE (corroded) and others. - SaR-32/S 1315; about 30 x.
- Fig. 4 ?*Nummulites fossulata* DE CIZANCOURT and others. - SaR-32/S 1315; about 30 x.
- Fig. 5 *Assilina laminosa* GILL, *Nummulites globulus* LEYMERIE, fragments of *Discocyclina* specimens and others. - SaR-32/S 1315; about 30 x.
- Fig. 6 *Assilina* aff. *laminosa* GILL and others. Note the reduced diameter and the increased thickness. - SaR-32/S 1315; about 30 x.

(a) the *Nummulites-Assilina pustulosa-Discocyclus rani-*
kotensis Assemblage of the Nammal Gorge section, western Salt
Range (Fig. 3),

(b) the *Nummulites-Assilina* Assemblage and probably the
Discocyclus-Assilina daviesi-Orbitolites complanatus Assem-
blage of the Panoba Anticline section, Kohat area (Fig. 6),

(c) the *Orbitolites complanatus-Discocyclus rani-*
kotensis-Nummulites fossilata Assemblage of the Mughal Kot-Toi
Nala section, Sulaiman Range (Fig. 7),

(d) the *Discocyclus-Nummulites globulus* Assemblage and
the *Alveolina-Nummulites fossilata* Assemblage of the Zinda
Pir section, Sulaiman Range (Fig. 8),

(e) the *Assilina leymeriei-Nummulites fossilata* Assemblage
of the Rakhi Nala section, Sulaiman Range (Fig. 10) and

(f) the *Nummulites globulus-Operculina salsa* Assemblage of
the Murree Brewery Gorge section (Fig. 11).

Apart from species modifications mentioned above the
assemblages are characterized by increased numbers of *Assilina*
specimens and by the appearance of the first *Nummulites*
specimens. These two well-known groups of larger forami-
nifers are represented mainly by abundant *Assilina leymeriei*
D'ARCHIAC & HAIME, *Nummulites globulus* LEYMERIE and
Nummulites atiacus LEYMERIE which occurs in younger parts
of the assemblage. Additional species are *Assilina laminosa*
GILL, *Assilina pustulosa* DONCIEUX and *Nummulites fossilata*
DE CIZANCOURT.

The assemblage is also characterized by a second abundance
maximum of *Discocyclus rani-*
kotensis DAVIES which was
found to be common in more fine-grained sediments, by
Orbitolites complanatus LAMARCK which occurs in more
coarse-grained sediments, and by abundant alveolinids (*Alveo-*
lina elliptica (SOWERBY), *Flosculina globosa* (LEYMERIE)) which
occur in quite shallow water environments. *Ranikothalia*
specimens have their last occurrence within basal parts of this
assemblage, *Discocyclus rani-*
kotensis DAVIES disappears at the
end of this assemblage, and *Nummulites fossilata* DE
CIZANCOURT is restricted to this assemblage.

The biostratigraphic age assignments with planktonic forami-
nifers as defined by TOUMARKINE & LUTERBACHER (1985) are
as follows:

(a) the *Nummulites-Assilina pustulosa-Discocyclus rani-*
kotensis Assemblage of the Nammal Gorge section (Fig. 3) was
found in higher parts of the Nammal Marls. Lower parts of the
Nammal Marls have been dated as belonging to the *Moro-*
zovella subbotinae and *Morozovella formosa* Zones of the
Early Eocene,

(b) the *Nummulites-Assilina* Assemblage of the Panoba
Anticline section (Fig. 6) occurs in the „Patala Shales“ (see
above) and was dated with co-occurring planktonic forami-
nifers as close to *Morozovella subbotinae* and *Morozovella*
formosa Zones of the Early Eocene. The *Discocyclus-Assilina*
daviesi-Orbitolites complanatus Assemblage which was iden-
tified from the Shekhan Formation of the Panoba Anticline
section (Fig. 6) is included here due to Early Eocene age
assignments by SHAH (1977: 759). This assemblage of larger
foraminifers is still uncertain biostratigraphically and some-
how transitional concerning the age assignment. In certain beds

of the Shekhan Limestone sequence of the Panoba Anticline
section (Fig. 6) there are rare, but remarkable *Discocyclus*
specimens which show a distinct sellate test morphology
similar to that of the Middle Eocene *Discocyclus sowerbyi*
NUTTALL. These specimens of *Discocyclus* might be the proba-
ble ancestor of *Discocyclus sowerbyi* NUTTALL.

(c) the *Orbitolites complanatus-Discocyclus rani-*
kotensis-Nummulites fossilata Assemblage of the Mughal Kot-Toi
Nala section (Fig. 7) occurs within the Ghazij Formation and
indicates by co-occurring planktonic foraminifers the *Moro-*
zovella subbotinae and *Morozovella formosa* Zones of the
Early Eocene,

(d) the *Discocyclus-Nummulites globulus* Assemblage from
the topmost part of the lower part of the Zinda Pir Limestone
and the *Alveolina-Nummulites fossilata* Assemblage of the
upper part of the Zinda Pir Limestone (Fig. 8) are of Early
Eocene age too as confirmed by the same planktonic forami-
nifer zones mentioned above,

(e) the *Assilina leymeriei-Nummulites fossilata* Assemblage
of the Rakhi Nala section (Fig. 10) was found within upper
parts of the Upper Rakhi Gaj Shales, the Green and Nodular
Shales and basal parts of the Rubbly Limestone. It could not
be dated exactly by means of planktonic foraminifers due to
the presence of long-ranging species. The age assignment as
Early Eocene is according to EAMES (1952) and NAGAPPA
(1959). Upper parts of the Upper Rakhi Gaj Shales were
assigned to planktonic foraminifer *Morozovella formosa*
Zone of the Early Eocene,

(f) the *Nummulites globulus-Operculina salsa* Assemblage of
the highest parts of the Paleogene Brewery Limestone (Fig. 11)
does not contain planktonic foraminifers. The Early Eocene
age assignment is according to ALLEMANN (1979).





Assilina spinosa-Flosculina globosa-Dictyoconoides cooki Assemblage

Age: late Early Eocene to early Middle Eocene

Remarks: The *Assilina spinosa-Flosculina globosa-Dictyo-*
conoides cooki Assemblage is intermediate between the *Assilina*
leymeriei-Nummulites fossilata-Discocyclus rani-
kotensis Assemblage of undoubtedly Early Eocene age (discussion above)
and the *Discocyclus dispansa-Discocyclus sowerbyi* Assem-
blage of undoubtedly Middle Eocene age (discussion below).

The assemblage of larger foraminifers is likewise a composite
characterized by a predominance either of *Assilina spinosa*
DAVIES and other assiliniids, such as *Assilina laciniata* DE
CIZANCOURT and *Assilina spinosa* DAVIES, which occur in more
coarse-grained and probably more turbulent shallow water
environments, or by a remarkable predominance of *Flosculina*
globosa (LEYMERIE) and *Alveolina elliptica* (SOWERBY) which
were found in more fine-grained and probably quite shallow
water or lagoonal environments. The environments may be
indicative of inner parts of the Sublittoral.

Dictyoconoides cooki (CARTER) seems to be restricted to this
interval. *Orbitolites complanatus* LAMARCK and *Nummulites*
specimens of both the *Nummulites globulus* and *Nummulites*
atacius group are present.



WEISS, W.: Maastrichtian to Eocene foraminiferal assemblages of Pakistan

Plate 3

- Plate 3** Nammal Gorge section, Salt Range. Larger and planktonic foraminifers.
- Fig. 1 Larger foraminiferal assemblage with *Discocyclina ranikotensis* DAVIES, *Nummulites globulus* LEYMERIE, *Nummulites* sp., *Assilina* sp. and others. Matrix with many planktonic foraminifers. - SaR-32/S 1315; about 8 x; Early Eocene.
- Fig. 2 *Discocyclina* sp. and others. Note the asymmetric thickness at the poles and the numerous polar bosses. - SaR-32/S 1315; about 30 x; Early Eocene.
- Fig. 3 *Nummulites globulus* LEYMERIE, *Rotalia trochidiformis* (LAMARCK) and others. - SaR-32/S 1315; about 30 x; Early Eocene.
- Fig. 4 *Nummulites ataticus* LEYMERIE, *Lockbartia* aff. *hamei* DAVIES and others. - SaR-32/S 1315; about 20 x; Early Eocene.
- Fig. 5 *Assilina* aff. *pustulosa* DONCIEUX, *Assilina lamnosa* GILL and others. - SaR-32/S 1315; about 30 x; Early Eocene.
- Fig. 6 *Morozovella* aff. *subbotinae* (MOROZOVA). - SaR-28/S 1377; about 60 x; Early Eocene.
- Fig. 7 *?Pseudobastigerina wilcoxensis* (CUSHMAN & PONTON). - SaR-28/S 1377; about 60 x; Early Eocene.
- Fig. 8 *Morozovella* aff. *gracilis* (BOLLI). - SaR-32/S 1315; about 60 x; Early Eocene.
- Fig. 9 *?Acarinina wilcoxensis* (CUSHMAN & PONTON). - SaR-32/S 1315; about 60 x; Early Eocene.

A more precise age of this assemblage was not verified by planktonic foraminifers. The *Assilina spinosa-Flosculina globosa-Dictyoconoides cooki* Assemblage was observed only from the Kohat Formation of the Panoba Anticline section, Kohat area (Fig. 6) which is of late Early Eocene to Middle Eocene age following SHAH (1977).

Discocyclina dispansa-Discocyclina sowerbyi Assemblage

Age: Middle Eocene

Remarks: This assemblage of larger foraminifers is well defined in age on the basis of Middle Eocene *Globigerinatheka subconglobata* and *Orbulinoides beckmanni* Zones of planktonic foraminifers and includes

(a) the *Planorbulina-Alveolina elliptica-Discocyclina dispansa* Assemblage of the Mughal Kot-Toi Nala section (Fig. 7),

(b) the *Discocyclina* Assemblage of the Zao River section (Fig. 9), and

(c) the *Discocyclina dispansa-Discocyclina sowerbyi* Assemblage of the Rakhi Nala section (Fig. 10).

It is characterized by the occurrence of *Discocyclina dispansa* (SOWERBY) and *Discocyclina sowerbyi* NUTTALL. They occur together with specimens of *Nummulites globulus* LEYMERIE and *Nummulites atacicus* LEYMERIE, *Operculina aspersa* COLOM, *Alveolina elliptica* (SOWERBY) and *Planorbulina* sp.

The assemblage is attributed to an open neritic, shallow-water environment of outer parts of the Sublittoral. It is widely distributed within the Pakistan-Indian bioprovince (SEN GUPTA 1963).

3.2.4 Remarks on Larger Foraminiferal Paleobiogeography

Paleobiogeographic considerations of Cretaceous and Paleogene larger foraminiferal assemblages from the Upper Indus Basin sections were not a main topic of investigations presented here. A comprehensive interpretation of biogeographic pattern of larger foraminifers requires basically detailed knowledge of type material of the abundant taxa of larger foraminifers which is essential to solve many problems of taxonomic interpretations still existant. Therefore, few aspects of larger foraminiferal paleobiogeography are given here as well as additional informations as they are related to the studied sections.

The importance of the shallow marine Paleogene sequences of Pakistan for the understanding of distribution of Paleogene larger foraminifers, and in particular that of the nummulitids and alveolins, is documented in many publications (cited already above and below) and is summarized for instance by BLONDEAU (1972) and KURESHI (1975, 1978). They accentuate that Paleogene shallow water carbonate deposits of Pakistan are essential of tropical and subtropical larger foraminiferal assemblages of the Far East and are a center of the Indo-Pakistan Province of the eastern Tethys region.


Paleocene and Eocene taxa of larger foraminifers from southern Tibet (WAN 1990) show close affinities to the species as described from Pakistan and emphasize the importance of the Indo-Pakistan region.

Larger foraminifers from the Baluchistan Basin, the Lower Indus Basin and the Upper Indus Basin of Pakistan were studied by many earlier paleontologists (SOWERBY 1837, D'ARCHIAC & HAIME 1853, CARTER 1861, NUTTALL 1926, DAVIES 1937, 1940, EAMES 1952, GILL 1953, SMOOTH & HAQUE 1956, NAGAPPA 1959, BAYLISS 1961, ADAMS 1970). Planktonic as well as larger foraminifers from Tertiary carbonate deposits of Pakistan were studied in detail by KURESHI (1966, 1969, 1971, 1972, 1974, 1975, 1978). This author has shown (and this is corroborated by the author's studies) that interregional stratigraphic correlation and paleobiogeographic comparison of larger foraminiferal assemblages is nevertheless complicated, even between the Paleogene deposits of the Baluchistan, Lower Indus and Upper Indus Basins, due to permanent problems of taxonomical interpretation of larger foraminifers and due to the restricted correlation of the stratigraphic ranges of larger foraminifers with planktonic foraminifers and other open marine microfossil groups. Comparison and correlation are also hampered because nowhere in Pakistan is a complete Tertiary carbonate sequence exposed in one stratigraphic section.

Concerning the geographic distribution of the Cretaceous orbitoidal foraminifers which „occupy a belt approximately parallel to present day low to low-middle latitudes in the Northern Hemisphere“ (VAN GORSEL 1978), this author has distinguished two main larger foraminiferal provinces on a generic level: a „Caribbean“ province of the western Tethys and a „Eurasian“ province of the eastern Tethys. Orbitoid genera *Omphalocyclus* and *Orbitoides* which were found in the Late Cretaceous limestones of the Sulaiman Range belong to world-wide genera and are widely distributed in America, Europe and into the Far East (VAN GORSEL 1978). *Omphalocyclus* and *Orbitoides* were not found in the Kohat area.

Compared with the rich Paleogene larger foraminiferal assemblages of the Lower Indus Basin (KURESHI 1975, 1978), it seems that the number of larger foraminiferal taxa of the Upper Indus Basin is reduced, if the number of *Nummulites*, *Assilina*, *Discocyclina* and *Alveolina* species described are compared.

If groups of larger foraminiferal assemblages of the Sulaiman Range, the Kohat area and the western Salt Range of the Upper Indus Basin are compared, there are some interregional differences (Fig. 12). Alveolins are rare in the Late Paleocene and Eocene sections of the western Salt Range. Nummulitids are more common in the Eocene intervals of the Sulaiman Range than in intervals of the western Salt Range and the Kohat area; *Nummulites fossulata* DE CIZANCOURT, *Nummulites crasseornata* (HENRICI) and *Nummulites vanderstoki* RUTTEN & VERMUNT were not found north of the Sulaiman Range which supports the assumption of a regional distribution pattern. Most of the Middle Eocene discocyclinids were found within the Sulaiman Range sections and were probably not able to immigrate and to occupy favourable paleoenvironments of the northern regions.



WEISS, W.: Maastrichtian to Eocene foraminiferal assemblages of Pakistan

Plate 4

Plate 4 Khairabad section, Salt Range. Algal and larger foraminiferal assemblages.

Fig. 1 Algal assemblage, mostly dasycladaceans and others. - SaR-59/S 1325; about 27 x; Middle Paleocene.

Fig. 2 *Miscellanea muscella* (D'ARCHIAC & HAIME) and others. - SaR-64/S 1331; about 30 x; Middle Paleocene.

Fig. 3 Large arenaceous specimen (triserial). - SaR-61/S 1328; about 30 x; Middle Paleocene.

Fig. 4 *Orbitolites complanatus* LAMARCA and others. - SaR-61/S 1328; about 30 x; Middle Paleocene.

Fig. 5 *Discocyclina ranikotensis* DAVIES (juvenile specimens) and others. Note the two-chambered nucleococonch. - SaR-60/S 1327; about 40 x; Middle Paleocene.

Fig. 6 *Discocyclina ranikotensis* DAVIES and others. Note the virtually three-chambered nucleococonch. - SaR-62/S 1329; about 40; Middle Paleocene.

4. SUMMARY

Late Cretaceous and Paleogene marine sediment sequences from the Upper Indus Basin of northern Pakistan (the western Salt Range, the Kohat area, the Sulaiman Range) were correlated on the basis of planktonic and larger foraminiferal assemblages. The studied sections are the Patala Nala section, the Nammal Gorge section, and the Khairabad section in the western Salt Range; the section North of Hangu and the Panoba Anticline section in the Kohat Area; and the Mughal Kot-Toi Nala section, the Zinda Pir section, the Zao river section, the Rakhi Nala section, and the Murree Brewery Gorge section in the Sulaiman Range. Cretaceous and Paleogene planktonic foraminifers of samples were used for comparison of various sections and areas as well as of different larger foraminiferal assemblages.

The assemblages of larger foraminifers identified are as follows:

- (1) the *Orbitoides media-Omphalocyclus macroporus* Assemblage of Early Maastrichtian age,
- (2) the *Siderolites calcitrapoides-Orbitoides media* Assemblage of early Late Maastrichtian age,
- (3) the *Siderolites calcitrapoides-Rotalia cf. trochidiformis* Assemblage of late Late Maastrichtian age,
- (4) the *Miscellanea-Lockhartia haimiei-Dictyokathina simplex* Assemblage of late Early to Middle Paleocene age,
- (5) the *Assilina dandotica-Discocyclina ranikotensis* Assemblage of Late Paleocene age,
- (6) the *Assilina leymyeri-Niommulites fosulata-Discocyclina ranikotensis* Assemblage of Early Eocene age,
- (7) the *Assilina spinosa-Flosculina globosa-Dictyoconoides cooki* Assemblage of late Early to early Middle Eocene age,

(8) the *Discocyclina dispansa-Discocyclina sowerbyi* Assemblage of Middle Eocene age.

Biostratigraphic analysis of microfaunas and in particular of planktonic and larger foraminifers revealed several first and last occurrences of planktonic and larger foraminifers in the sections of the western Salt Range, the Kohat area and the Sulaiman Range. They were used to refine regional stratigraphy.

The remarkable facies change from the pelagic Parh Formation to the flyschoid Mughal Kot Formation in the Mughal Kot-Toi Nala section was dated with the last occurrence of high-spired praeglobotruncanids and with the range of *Margmotruncana pseudolimeiana* as Late Turonian to Early Coniacian.

Stepwise regression during the Maastrichtian is documented in the Rakhi Nala section of the Sulaiman Range by the succession of different assemblages of larger foraminifers which indicate successively shallower environments within the *Orbitoides* Limestone Formation and by the facies change into the Pab Sandstone sequence.

Paleoenvironmental changes during the Paleogene have influenced the larger foraminiferal succession in particular at the Middle-Late Palaeocene boundary and the Palaeocene-Eocene boundary in the western Salt Range as well as in the Sulaiman Range. The shallow water carbonate facies ends in the western Salt Range near the end of the Early Eocene, in the Kohat area within the Middle Eocene and in the Sulaiman Range in the Late Eocene.

REFERENCES

- ADAMS, C. G. (1967): Tertiary foraminifera in the Tethys, America and Indo-Pacific Province. - In: ADAMS, C. G. & ÅGER, D. V. (eds.), Aspects of Tethyan Biogeography; Systematic Assoc. London, 195-217; London.
- ADAMS, C. G. (1970): A reconsideration of the East Indian Letter Classification of the Tertiary. - Brit. Mus. (Nat. Hist.) Bull. Geol., 19 (3): 87-137; London.
- ADAMS, C. G. (1989): Foraminifera as indicators of geological events. - Proc. Geol. Ass., 100 (3): 297-311; London/Edinburgh.
- ALLEMANN, F. (1979): Time of emplacement of the Zhob Valley ophiolites and Bela ophiolites, Baluchistan (Preliminary report). - In: FARAH, A. & DE JONG, K. A. (eds.), Geodynamics of Pakistan, 215-242; Quetta (Geol. Surv. Pakistan).
- ARCHIAC, E. J. A. DE (1837): Mémoire sur la formation crétacée du SW de la France. - Mém. Soc. géol. France, (2), 2: 157-192; Paris.
- ARCHIAC, E. J. A. DE & HAIME, J. (1853/1854): Description des animaux fossils du groupe Nummulitique de l'Inde. Précédé d'un résumé géologique et d'une monographie des Nummulites. - 2 vols., 373 pp.; Paris (Gide et Baudry).
- BANNERT, D., in cooperation with AMJED CHEEMA, ABRAR AHMED, and U. SCHÄFFER (1992): The structural development of the Western Fold Belt, Pakistan. - Geol. Jb., B 80: 3-60; Hannover.
- BAYLISS, B. B. (1961): An investigation of certain larger fossil foraminifera from Pakistan. - Ph. D. Thesis, Dept. Geol. Univ. Wales, 253 pp.; Cardiff.
- BLONDEAU, A. (1972): Les Nummulites. - 254 pp.; Paris (Vuibert).
- BLOW, W. A. (1969): Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. - In: BRÖNNIMANN, P. & RENZ, H. H. (eds.), Proc. 1. Intern. Conf. Plankt. Microfossils, Geneva 1967, 1: 199-422; Leiden (Brill).
- BLOW, W. A. (1979): The Cainozoic Globigerinida. A study of the morphology, taxonomy, evolutionary relationships and the stratigraphical distribution of some Globigerinida (mainly Globigerinacea). - 3 vols., 15 + 1413 pp.; Leiden (Brill).
- BOLLI, H. M. (1957): The genera *Praeglobotruncana*, *Rotalipora*, *Globotruncana*, and *Abatbomphalus* in the Upper Cretaceous of Trinidad, B. W. I. - Bull. U. S. Nat. Mus., 215: 51-60; Washington.
- BOLLI, H. M. (1966): Zonation of Cretaceous to Pliocene marine sediments based on planktonic foraminifera. - Bol. Inf. Ass. Venezolana Geol., Min. and Petrol., 9 (1): 3-33; Chacao.
- BOLTOVSKOY, E. & WRIGHT, R. (1976): Recent Foraminifera. - 515 pp.; Den Haag (Junk).
- BUTT, A. A. (1987): The Paleogene stratigraphy of the Kala Chitta Range, Northern Pakistan. - Acta Min. Pakistan., 3: 97-110.




Plate 5





Plate 5 Khairabad section, Salt Range. Larger foraminifera.

- Fig. 1 Larger foraminiferal assemblage with *Orbitolites complanatus* LAMARCK, *Operculina salsa* DAVIES and others. - SaR-3/S 1309; about 6 x; Middle Paleocene.
- Fig. 2 *Orbitolites complanatus* LAMARCK, *Dictyokathina simplex* SMOUTH and others. - SaR-3/S 1309; about 6 x; Middle Paleocene.
- Fig. 3 *Lockhartia conditi* (NUTTALL), *Kathina* aff. *selverti* SMOUTH and others. - SaR-3/S 1309; about 40 x; Middle Paleocene.
- Fig. 4 *Miscellanea miscella* (D'ARCHIAC & HAIME), ?*Dictyokathina simplex* (juvenile specimen) and others. - SaR-3/S 1309; about 30 x; Middle Paleocene.
- Fig. 5 *Operculina salsa* DAVIES, *Miscellanea miscella* (D'ARCHIAC & HAIME) and others. - SaR-3/S 1309; about 23 x; Middle Paleocene.
- Fig. 6 *Miscellanea miscella* (D'ARCHIAC & HAIME) and others. - SaR-3/S 1309; about 30 x; Middle Paleocene.

- BLITT, A. A. (1991): *Ranikothalia sindensis* zone in Late Paleocene biostratigraphy. - *Micropaleontology*, 37 (1): 77-85; New York.
- BUTTERLIN, J., FOURCADE, E. (1989): Stratigraphic range and geographic distribution of the genus *Lockhartia* DAVIES, 1932 (Foraminifera, Rotaliidae). - *Rev. Micropal.*, 31 (4): 225-242; Paris.
- CARON, M. (1985): Cretaceous planktic foraminifera. - In: BOLLI, H. M., SAUNDERS, J. B. & PERCH-NIELSEN, K. (eds.), *Plankton stratigraphy*, 17-86; Cambridge (Cambridge Univ. Press).
- CARTER, H. J. (1861): Further observations on the structure of Foraminifera and on larger fossilized forms of Sind, etc., including a new genus and species. - *Ann. Mag. Nat. Hist. London*, (3), 8 : 246-251, 309-333, 366-382, 446-470; London.
- CIZANCOURT, H. DE (1938): Nummulites et Assilines du flysch de Gardez et du Khost, Afghanistan oriental. - In: CIZANCOURT, A. DE & COX, L. R.: *Contribution à l'étude des faunes tertiaires de l'Afghanistan*. - *Mém. Soc. Géol. France*, N. S., 39: 5-28; Paris.
- CIZANCOURT, H. DE (1946): Dénomination nouvelles pour *Assilina umblicata* DE CIZANCOURT, 1938. - *C. R. Soc. géol. France*, 16: 335; Paris.
- COLOM, G. (1954): Estudio de las biozonas con foraminíferos del Terciario de Alicante. - *Bol. Inst. Geol. Min. Espan.*, 66: 183; Madrid.
- CONCLUSIONS/NICE (1983): Conclusions au Colloque sur les Etages Coniacien à Maastrichtien: Echelles biostratigraphiques. - *Géol. Méditerran.*, 10 (3/4): 413-434; Nice.
- DAVIES, L. M. (1926): Remarks on CARTER'S genus *Conulites-Dictyoconoides* NUTTALL, with descriptions of some species from the Eocene of North West India. - *Rec. Geol. Surv. India*, 59: 237-253; Calcutta.
- DAVIES, L. M. (1927): The Ranikot Beds at Thal (North West Frontier Provinces of India). - *Quart. Jour. Geol. Soc. London*, 83 (1): 260-290; London.
- DAVIES, L. M. (1932): The genera *Dictyoconoides* NUTTALL, *Lockhartia* nov. and *Rotalia* LAMARCK. - *Trans. Roy. Soc.*, 57: 397-428; Edinburgh.
- DAVIES, L. M. (1940): The Upper Kirthar Beds of North West India. - *Quart. Jour. Geol. Soc. London*, 96 (2): 199-230; London.
- DAVIES, L. M. & PINFOLD, E. S. (1937): The Eocene Beds of the Punjab Salt Range. - *Mem. Geol. Surv. India, Palaeontologica India*, N. S., 24 (1): 14-79; Calcutta.
- DONCIEUX, L. (1926): Catalogue descriptif des fossiles Nummulitiques de l'Aude et de l'Hérault. - *Ann. Univ.*, n. s., 45: 1-99; Lyon.
- DOUVILLE, H. (1919): L'Eocène inférieur en Aquitaine et dans les Pyrénées. - *Mém. Explic. Carte géol. dét. France*, 1-84; Paris.
- EAMES, F. E. (1952a): A contribution to the study of the Eocene in Western Pakistan and Western India: A. The geology of standard sections in the Western Punjab and in the Kohat district. - *Quart. Jour. Geol. Soc. London*, 107: 150-171; London.
- EAMES, F. E. (1952b): A contribution to the study of the Eocene in Western Pakistan and Western India. D. Discussion of the faunas of certain standard sections, and their bearing on the classification and correlation of the Eocene in Western Pakistan and Western India. - *Quart. Jour. Geol. Soc. London*, 107: 173-200; London.
- FATMI, A. N. (1973): Lithostratigraphic units of the Kohat Potwar Province, Indus Basin, Pakistan. - *Mem. Geol. Surv. Pakistan*, 10: 80 pp.; Islamabad.
- FATMI, A. N. & KHAN, M. R. (1966): Stratigraphy of parts of western Kohat, West Pakistan (Samana-Darsamand and Thal sections). - *Rec. Geol. Surv. Pakistan*, 20; Quetta.
- FRITZ, E. B. & KHAN, M. R. (1967): Cretaceous (Albian-Cenomanian) planktonic Foraminifera in Bangu Nala, Quetta Division, West Pakistan. - *U. S. Geol. Surv. Proj. Rep.*, PK-36: 16 pp; Washington.
- GEE, E. R. (1980): Pakistan geological Salt Range map series (6 sheets, scale 1: 50 000). - Directorate of Overseas Surveys, United Kingdom, for the Government of Pakistan and Geological Survey of Pakistan.
- GEE, E. R. (1989): Overview of the geology and structure of the Salt Range, with observations on related areas of northern Pakistan. - In: MALINCONICO, L. L. & LILLIE, R. J. (eds.), *Tectonics of western Himalayas*, *Spec. Pap. Geol. Soc. Amer.*, 232: 95-112; Boulder.
- GIGON, W. O. (1962): Upper Cretaceous stratigraphy of the well Giandari-1 and its correlation with the Sulaiman and Kirthar Ranges, West Pakistan. - *ECAFE Symp. Div. Petrol. Res. Asia and Far East*, 282-284; Teheran.
- GILL, W. D. (1952): Facies and fauna in the Bhadrar Beds of the Punjab Salt Range, Pakistan. - *Jour. Paleont.*, 27 (6): 824-844; Menasha.
- GILL, W. D. (1953): The genus *Assilina* in the Laki Series (Lower Eocene) of the Kohat-Potwar Basin, Northwest Pakistan. - *Contrib. Cush. Found. For. Res.*, 4 (2): 76-84; Washington.
- GORSSEL, J. T. VAN (1970): Late Cretaceous orbitoidal Foraminifera. - In: HEDLEY, R. H. & ADAMS, C. G. (eds.), *Foraminifera*, 1-120; London (Academic Press).
- HAAR, R. & POSTUMA, J. (1975): The relation between tropical planktonic foraminiferal zonation and the Tertiary Far East Letter Classification. - *Geologie en Mijnbouw*, 54 (3-4): 195-198; Den Haag.
- HAQUE, A. F. M. M. (1956): The smaller Foraminifera of the Ranikot and the Laki of the Nammal Gorge, Salt Range. - *Mem. Geol. Surv. Pakistan*, 1: 1-300.
- HENRICI, H. (1934): Foraminiferen aus dem Eozän und Altmiozän von Timor. - In: BOEHM, G. & WANNER, J., *Beiträge zur Geologie von Niederländisch-Indien*, (4), 4 (1): 32, 52; Stuttgart.
- HOEKER, J. (1926): Die Foraminiferen aus dem Senon Limburgens. I: *Calcarina calcirapoides* LAMARCK. - *Naturhist. Maandbl.*, 15 (2): 14-17; Maastricht.
- HOEKER, J. (1963): Foraminifera from the Cretaceous of South-Limburg, Netherlands. 67: The taxonomic position of *Siderolites calcirapoides* LAMARCK. - *Naturhist. Maandbl.*, 52 (7-8): 109-114; Maastricht.
- HOTTINGER, L. (1960): Recherches sur les Alvéolines du Paléocène et de l'Eocène. - *Schweiz. paläont. Abh.*, 75/76: 144 pp.; Basel.

Plate 6 Khairabad section, Salt Range. Larger foraminifers.




- Fig. 1 Larger foraminiferal assemblage. - SaR-66/S 1334; about 6 x; Early Eocene.
- Fig. 2 Larger foraminiferal assemblage with *Discocyclina ranikotensis* DAVIES, *Nummulites* aff. *atacius* LEYMERIE and others. - SaR-66/S 1334; about 10 x; Early Eocene.
- Fig. 3 *Nummulites* aff. *atacius* LEYMERIE and others. - SaR-66/S 1334; about 10 x; Early Eocene.
- Fig. 4 ?*Assilina* aff. *plana* SCHAUB and others. - SaR-66/S 1334; about 15 x; Early Eocene.
- Fig. 5 *Nummulites* aff. *atacius* LEYMERIE (juvenile specimen) and others. - SaR-66/S 1334; about 40 x; Early Eocene.
- Fig. 6 *Assilina* aff. *leymeriei* D'ARCHIAC & HAIME and others. - SaR-66/S 1334; about 30 x; Early Eocene.
- Fig. 7 Equatorial sections of ?*Assilina* aff. *plana* SCHAUB (questionable A generation) and *Discocyclina ranikotensis* DAVIES. - SaR-65a/S 1333; about 15 x; Early Eocene



- HOTTINGER, L. (1960): Über paleocäne und eocäne Alveolinen. - *Eclogae geol. Helvet.*, 53 (1): 265-283; Basel.
- HOTTINGER, L. (1971): Larger Foraminifera common to Mediterranean and Indian Paleocene and Eocene formations. - *Ann. Inst. Geol. Publ.*, 54 (4/1): 145-151; Budapest.
- HUNTING SURVEY CORPORATION (1961): Reconnaissance geology of part of West Pakistan (Colombo Plan Cooperative Project). - 555 pp.; Toronto.
- JAMIL AFZAL & VON DANIELS, C. H. (1991): Foraminiferal biostratigraphy and paleoenvironment interpretation of the Paleocene to Eocene Patala and Nammal Formations from Khairabad-East, western Salt Range, Pakistan. - *Pak. Jour. Hydrocarbon Res.*, 3 (2): 61-79; Islamabad.
- KAEVER, M. (1964): Über *Assilina lacumata* CIZANCOURT aus dem Unter-Eozän Afghanistans. - *Geol. Jb.*, 82: 131-142; Hannover.
- KÖTHE, A. (1987): Jurassic to Paleogene nannofossil and Paleogene dinoflagellate biostratigraphy of the Surghar Range, Salt Range, Sulaiman Range and the Kohat area, Pakistan. - BGR Report: 86 pp.; Hannover.
- KÖTHE, A. (1988) in cooperation with A. M. KHAN & M. ASHRAF: Biostratigraphy of the Surghar Range, Sulaiman Range and Kohat area, Pakistan, according to Jurassic through Paleogene calcareous nannofossils and Paleogene dinoflagellates. - *Geol. Jb.*, B 71: 3-87; Hannover.
- KUPPER, K. (1954): Notes on the Upper Cretaceous larger Foraminifera. II: Genera of the subfamily Orbitoidinae with remarks on the microspheric generation of *Orbitoides* and *Omphalocyclus*. - *Contr. Cushman Found. Foram. Res.*, 5 (4): 179-184; Washington.
- KURESHY, A. A. (1966): Biostratigraphic studies of Chharat and adjacent areas of West Pakistan. - *Bull. Coll. Sci.*, 9: 85-92; Baghdad.
- KURESHY, A. A. (1969): Eocene biostratigraphy of West Pakistan. - *Mém. BRGM*, 69 (3): 219-224; Orléans.
- KURESHY, A. A. (1971): The Upper Cretaceous and Paleocene larger Foraminifera of Murree Brewery, West Pakistan. - *Proc. 4. African Micropal. Coll.*, 220-227.
- KURESHY, A. A. (1972): Stratigraphic micropaleontology of Pakistan. - *Pakistan Jour. Sci.*, 24 (3): 133-146; Lahore.
- KURESHY, A. A. (1974): The biostratigraphic correlation of sedimentary basins of Pakistan. - *Proc. 6. African Micropal. Coll.*, 327-336; Tunis.
- KURESHY, A. A. (1975): The Cretaceous and Tertiary larger Foraminifera of West Pakistan. - *Rev. Espan. Micropaleont.*, spec. publ., 7: 553-564; Madrid.
- KURESHY, A. A. (1975/1977): Taxonomical studies of Tertiary larger foraminifera. - *Rec. Geol. Surv. Pakistan*, 38: 1-47; Karachi.
- KURESHY, A. A. (1976): The Cretaceous planktonic foraminiferal zones of Pakistan. - *Rev. Espan. Micropaleont.*, 8 (3): 429-438; Madrid.
- KURESHY, A. A. (1977): The Tertiary planktonic foraminiferal zones of Pakistan. - *Rev. Espan. Micropaleont.*, 9 (2): 203-219; Madrid.
- KURESHY, A. A. (1978): Tertiary larger foraminiferal zones of Pakistan. - *Rev. Esp. Micropaleont.*, 10 (3): 467-483; Madrid.
- KURESHY, A. A. (1980): Larger foraminiferal biostratigraphy of the Kirthar Formation of Pakistan. - *Geology and Palaeontology of Southeast Asia*, 21: 235-245; Tokyo.
- KURESHY, A. A. (1984): The foraminiferal stratigraphy of Laki Formation (Early Eocene) of Pakistan. - *Géol. Méditerran.*, 11 (2): 231-236; Nice.
- LAMARCK, J. B. DE (1801): *Système des animaux sans vertèbres*. - 8+432 pp.; Paris.
- LAMARCK, J. B. DE (1816): *Histoire naturelle des animaux sans vertèbres*. - Paris.
- LATIF, M. A. (1961): The use of pelagic Foraminifera in the subdivision of the Paleocene-Eocene of the Rakhi Nala, West Pakistan. - *Geol. Bull. Panjab Univ.*, 1: 31-46; Lahore.
- LATIF, M. A. (1963): Some related groups of pelagic Foraminifera in the Paleocene-Eocene of the Rakhi Nala, West Pakistan. - *Geol. Bull. Panjab Univ.*, 3: 19-24; Lahore.
- LATIF, M. A. (1964): Variations in abundance and morphology of pelagic Foraminifera in the Paleocene-Eocene of the Rakhi Nala, West Pakistan. - *Geol. Bull. Panjab Univ.*, 4: 29-100; Lahore.
- LATIF, M. A. (1970): Explanatory notes on the geology of South-Eastern Hazara, to accompany the revised geological map. - *Jb. Geol. B.-A., Sonderbd.*, 15: 5-19; Wien.
- LATIF, M. A. (1970): Micropaleontology of the Chanali Limestone, Upper Cretaceous of Hazara, West Pakistan. - *Jb. Geol. B.-A., Sonderbd.*, 15: 25-61; Wien.
- LATIF, M. A. (1970): Micropaleontology of the Galis Group, Hazara, West Pakistan. - *Jb. Geol. B.-A., Sonderbd.*, 15: 63-66; Wien.
- LEYMERIE, A. (1846): Mémoire sur le terrain à *Nummulites* (épéricrécé) des Corbières et de la Montagne Noir. - *Mém. Soc. géol. France*, (2), 1: 337-373; Paris.
- MARKS, P. (1962): Variation and evolution in *Orbitoides* of the Cretaceous of the Rakhi Nala, West Pakistan. - *Bull. Geol. Univ. Panjab*, 2: 15-24; Lahore.
- MEISSNER, CH. R. et al. (1974): Stratigraphy of the Kohat Quadrangle, Pakistan. - *U. S. Geol. Surv. Prof. Pap.*, 716-D: 1-29; Washington.
- NAGAPPA, Y. (1959): Foraminiferal biostratigraphy of the Cretaceous-Eocene succession in the India-Pakistan-Burma region. - *Micropaleontology*, 5 (2): 145-192; New York.
- NASER A. SHAFIQUE & VON DANIELS, C. H. (1990): Foraminiferal Zonation of Upper Goru Formation - Bawani Area, Kirthar Range. - *Pak. Jour. Hydrocarbon Res.*, 2 (2): 67-84; Islamabad.
- NEUMANN, M. (1958): Révision des Orbitoididés du Crétacé et de l'Eocène en Aquitaine Occidentale. - *Mém. Soc. géol. France*, n. s., 37 (2-3), 83: 1-174; Paris.
- NICORA, A., GARZANTI, E., FOIS, E. (1986): Evolution of the Tethys Himalaya continental shelf during Maastrichtian to Paleocene (Zanskar, India). - *Riv. Ital. Paleont. Strat.*, 92 (4): 439-496; Milano.
- NUTTALL, W. L. F. (1925a): The stratigraphy of the Laki Series (Lower Eocene) of parts of Sind and Baluchistan (India); with a description of the larger Foraminifera contained in those beds. - *Quart. Jour. geol. Soc.*, 81 (3): 417-453; London.
- NUTTALL, W. L. F. (1925b): Two species of Eocene Foraminifera from India. *Alveolina elliptica* and *Dictyoconoides cooki*. - *Ann. Mag. Nat. Hist.*, (9), 16: 378-388; London/Edinburgh.

Plate 7 North of Hangu section, Kohat Area. Larger foraminifers.

- Fig. 1 Algal-foraminiferal assemblage. - Ko-59/S 1440; about 60 x; Late Maastrichtian.
- Fig. 2 Algal assemblage, mostly dasycladaceans, and larger foraminifers. - Ko-61/S 1442; about 20 x; ?Early to Middle Paleocene.
- Fig. 3 *Rotalia* cf. *trochidiformis* (LAMARCK), *Gublerina* sp. (left) and others. - Ko-59/S 1440; about 60 x; Late Maastrichtian.
- Fig. 4 *Rotalia trochidiformis* (LAMARCK), *Kathina major* SMOOTH and others. - Ko-61/S 1442; about 30 x; ?Early to Middle Paleocene.
- Fig. 5 Questionable soritid specimens. - Ko-59/S 1440; about 46 x; Late Maastrichtian.
- Fig. 6 Transitional specimen between *Lockhartia bamei* (DAVIES) and *Lockhartia atispina* SMOOTH and others. - Ko-61/S 1442; about 40; ?Early to Middle Paleocene.
- Fig. 7 *Rotalia* cf. *trochidiformis* (LAMARCK) (horizontal section). Note the thin, laminated, radially fibrous calcite of the outer wall. - Ko-59/S 1440; about 40; Late Maastrichtian.



- NUTTALL, W. L. F. (1926a): The larger Foraminifera of the Upper Ranikot Series (Lower Eocene) of Sind, India. - *Geol. Magazine*, 63 (3): 112-121; London.
- NUTTALL, W. L. F. (1926b): The zonal distribution of larger Foraminifera of the Eocene of Western India. - *Geol. Magazine*, 63 (11): 495-504; London.
- ORBIGNY, A. DE (1826): Tableau méthodique de la classe des Céphalopodes. - *Ann. Sci. Nat. Paris*, (1), 7: 96-314; Paris.
- OVEY, C. D. (1947): A new Eocene species of *Loekbarta* DAVIES, from British Somaliland, with notes on other species of the genus. - *Ann. Mag. Nat. Hist.*, (2), 13: 571-576; London.
- PALMER, D. K. (1934): Some large fossil foraminifera from Cuba. - *Mem. Soc. Cubana Nat. Hist.*, 8 (4): 235-264; Cuba.
- PAPP, A. & KUPPER, K. (1953): Die Foraminiferenfauna von Guttaring und Klein St.-Paul. II: Orbitoiden aus Sandsteinen vom Pemberg bei Klein St.-Paul. III: Foraminifera aus dem Campan von Silberberg. - *Sitzber. Akad. Wiss. Wien, Abt. I*, 162: 65-82; 345-357; Wien.
- PFENDER, J. (1935): A propos du *Siderolites vidali* DOUVILLÉ et de quelques autres. - *Bull. Soc. géol. France*, (5), 4: 225-236; Paris.
- PLUMMER, H. (1931): Some Cretaceous Foraminifera in Texas. - *Univ. Texas Bull.*, 3101: 109-237; Austin.
- PORTH, H., HILAL A. RAZA (1990a): On the geology and hydrocarbon prospects of Potwar Depression and adjoining sedimentary areas. - *BGR/HDIP Report*: 193 pp.; Hannover.
- PORTH, H., HILAL A. RAZA (1990b): On the geology and hydrocarbon prospects of Sulaiman Province, Indus Basin, Pakistan. - *BGR/HDIP Report*: 127 pp.; Hannover.
- POSTUMA, J. (1971): *Manual of planktonic Foraminifera*. - 420 pp.; Amsterdam (Elsevier).
- REICHEL, M. (1950): Observations sur les *Globotruncana* du gisement de la Breggia (Tessin). - *Eclogae geol. Helv.*, 42 (2): 596-617; Basel.
- ROBASZYNSKI, F. & CARON, M. (eds.) (1979): *Atlas de Foraminifères planctoniques du Crétacé moyen (Mer boréale et Téthys)*. Parts 1-2. - *Cah. Micropaléont.*, 1: 1-185; 2: 1-181; Paris.
- ROBASZYNSKI, F., CARON, M., GONZALES, J. M. & WONDERS, A. (eds.) (1984): *Atlas of Late Cretaceous globotruncanids*. - *Rev. Micropaléont.*, 26 (3-4): 145-305; Paris.
- RUTTEN, M. G. & VERMUNT, L. W. J. (1932): The Serce di Cueba limestone from Curacao. - *Proc. K. Akad. Wetensch.*, 35: 240; Amsterdam.
- SAMANTA, B. K. (1972): Planktonic foraminiferal biostratigraphy of the Early Tertiary of the Rakhi Nala section, Sulaiman Range, Pakistan. - *Jour. Geol. Soc. India*, 13 (4): 317-328; Bangalore.
- SAMANTA, B. K. (1973): Planktonic Foraminifera from the Palaeocene-Eocene succession in the Rakhi Nala, Sulaiman Range, Pakistan. - *Bull. Brit. Mus. (Nat. Hist.)*, 22 (6): 421-482; London.
- SCHAUB, H. (1981): Nummulites et Assilines de la Téthys paléogène. Taxinomie, phylogénèse et biostratigraphie. - *Schweiz. Paläontol. Abh.*, 104: 236 p.; 105 (Atlas 1): pls. 1-48; 106 (Atlas 2): pls. 49-97; Basel.
- SCHLUMBERGER, C. (1901): Première note sur les *Orbitoides*. - *Bull. Soc. géol. France*, (4), 1: 459-467; Paris.
- SCHLUMBERGER, C. (1902): Deuxième note sur les *Orbitoides*. - *Bull. Soc. géol. France*, (4), 2: 255-261; Paris.
- SCHWAGER, C. (1883): Die Foraminiferen aus den Eocaenablagerungen der Libyenschen Wüste und Aegyptens. - *Palaeontographica*, (3/6), 30: 79-153; Cassel.
- SEN GUPTA, B. K. (1963): A restudy of two common species of *Discocyclina* from India. - *Micropalaeontology*, 9 (1): 39-49; New York.
- SHAH, S. M. J. (1977): Stratigraphy of Pakistan. - *Mem. Geol. Surv. Pakistan*, 12: 1-138; Islamabad.
- SIDDIQUI, Q. A. (1971): Early Tertiary Ostracoda of the family Trachyleberididae from West Pakistan. - *Bull. Brit. Mus. (Nat. Hist.)*, 9: 1-99; London.
- SMOUTH, A. H. (1954): Lower Tertiary Foraminifera from the Qatar Peninsula. - 96 pp.; London.
- SMOUTH, A. H. & HAQUE, A. F. M. M. (1956): A note on the larger Foraminifera and Ostracoda of the Ranikot from the Namnal Gorge, Salt Range, West Pakistan. - *Rec. Geol. Surv. Pakistan*, 8 (2): 49-60; Karachi.
- SOWERBY, J. DE C. (1840): Systematic list of organic remains. Appendix to GRANT, C. W., Memoir to illustrate a geological map of Cutch. - *Trans. Geol. Soc. London*, (2), 5: 327-329; London.
- STAINFORTH, R. M., LAMB, J. L., LUTERBACHER, H., BEARD, J. H., JEFFORDS, R. M. (1975): Cenozoic planktonic foraminiferal zonation and characteristics of index forms. - *Paleontol. Contrib. Univ. Kansas*, 62: 425 pp.; Lawrence.
- TOUMARKINE, M. & LUTERBACHER, H. (1985): Palaeocene and Eocene planktonic foraminifera. - In: BOLLI, H. M., SAUNDERS, J. B. & PERCH-NIELSEN, K. (eds.): *Plankton stratigraphy*, 87-154; Cambridge (Cambridge Univ. Press).
- WAN, X. (1990): Eocene larger Foraminifera from Southern Tibet. - *Rev. Espan. Micropaléont.*, 22 (1): 213-238; Madrid.
- WEISS, W. (1988): Larger and planktonic foraminiferal biostratigraphy of the Cretaceous and the Paleogene in the Salt Range, the Kohat area and the Sulaiman Range, Pakistan. - *BGR Report*: 57 pp.; Hannover.
- WILLIAMS, M. D. (1959): Stratigraphy of the Lower Indus Basin, West Pakistan. - *5th World Petrol. Congr. Proc.*, (1), 19: 377-390; New York.

APPENDIX

Faunal reference list of larger foraminifers

Taxa of larger foraminifers identified in the samples are listed alphabetically with reference to their original description. Additional references of preferred modern nomenclature of larger foraminifers (including as far as possible A and B generations) are given showing typical species illustrations which were the base of determinations used here. Some taxa in open nomenclature are discussed.

Cretaceous

Omphalocyclus macroporus (LAMARCK, 1816)

= *Orbulites macropora* n. sp. - LAMARCK: 197.

= *Omphalocyclus macroporus* (LAMARCK) - NEUMANN (1958): 65-66; pl. 6, figs. 1-8; pl. 35, fig. 2; text-fig. 16a-b.

Orbitoides apiculata SCHLUMBERGER, 1901

= *Orbitoides apiculata* n. sp. - SCHLUMBERGER: 462; pl. 8, figs. 1, 4, 6; pl. 9, figs. 1, 4.


= *Orbitoides apiculata apiculata* (SCHLUMBERGER) - KUPPER (1954): 66-67; pl. 12, figs. 7-8;

= *Orbitoides apiculata* (SCHLUMBERGER) - NEUMANN (1958): 63-65; pl. 4, figs. 1-6; pl. 5, figs. 5-6; text-fig. 15a-b.

Orbitoides media (D'ARCHIAC, 1837)

= *Orbitolites media* n. sp. - D'ARCHIAC: 178;

= *Orbitoides media media* (D'ARCHIAC) - PAPP & KUPPER (1953): 73; pl. 1, figs. 5-7; pl. 2, figs. 2-4;







WEISS, W.: Maastrichtian to Eocene foraminiferal assemblages of Pakistan

Plate 8

- Plate 8** Rakhi Nala section, Sulaiman Range. Larger foraminifers.
- Fig. 1 *Orbitoides tissoti* SCHLUMBERGER. Note the different sections probably indicating a low-energy depositional environment. - SmR-1/S 1419; about 6 x; Early Maastrichtian.
 - Fig. 2 *Orbitoides tissoti* SCHLUMBERGER. Note the predominant axial sections probably indicating a high-energy depositional environment. - SmR-1/S 1419; about 8 x; Early Maastrichtian.
 - Fig. 3 *Orbitoides tissoti* SCHLUMBERGER. Note the two and three tubelike stolons serving as a connection between chambers of the central layer. - SmR-1/S 1419; about 46 x; Early Maastrichtian.
 - Fig. 4 *Orbitoides tissoti* SCHLUMBERGER. Note the virtually three-chambered nucleocoench. - SmR-1/S 1419; about 40 x; Early Maastrichtian.
 - Fig. 5 *Orbitoides tissoti* SCHLUMBERGER. Note the quadrilocular nucleocoench surrounded by thick, perforated wall. - SmR-1/S 1419; about 46 x; Early Maastrichtian.
 - Fig. 6 *Orbitoides tissoti* SCHLUMBERGER. Note the arcuate equatorial chambers in a slightly oblique section. - SmR-1/S 1419; about 23 x; Early Maastrichtian.

- = *Orbitoides media* (D'ARCHIAC) - NEUMANN (1958): 60-62; pl. 3, figs. 1-6; pl. 5, figs. 3-4; pl. 35, fig. 2; text-fig. 14a-d.
- Orbitoides tissoti* SCHLUMBERGER, 1902
- = *Orbitoides tissoti* n. sp. - SCHLUMBERGER: 259; pl. 8, figs. 21-25.
- = *Orbitoides tissoti tissoti* (SCHLUMBERGER) - PAPP & KUPPER (1953): 349; pl. 1, figs. 1-2; pl. fig. 2;
- = *Orbitoides tissoti* (SCHLUMBERGER) - NEUMANN (1958): 56-60; pl. 1, figs. 1-9; pl. 2, figs. 1-4; pl. 5, figs. 1-2; pl. 35, fig. 1; text-fig. 13a-b.
- Siderolites calcitrapoides* LAMARCK, 1801
- = *Calcarina calcitrapoides* (LAMARCK) - HOFKER (1926): 14-17; figs. 1-14.
- Paleogene**
- Alveolina delicatissima* SMOUTH, 1954
- = *Alveolina delicatissima* n. sp. - SMOUTH: 83-84; pl. 14, fig. 13.
- Alveolina elliptica* (SOWERBY, 1840)
- = *Fasciolites elliptica* n. sp. - SOWERBY: 329; pl. 24, fig. 17;
- = *Alveolina elliptica* (SOWERBY) - HOTTINGER (1960): 146; pl. 12, figs. 1-3.
- Alveolina lepidula* SCHWAGER, 1883
- = *Alveolina lepidula* var. A - SCHWAGER: pl. 25, figs. 3a-g;
- = *Alveolina lepidula* var. A - SCHWAGER - NUTTALL (1925): 439; pl. 24, figs. 1-2;
- = *Alveolina* (*Glomalveolina*) *lepidula* SCHWAGER - HOTTINGER (1960): 57-58; pl. 1, figs. 25-29; pl. 2, figs. 9, 25; text-figs. 25, 29: 20-22, 80.
- Alveolina oblonga* D'ORBIGNY, 1826
- = *Alveolina oblonga* n. sp. - D'ORBIGNY: 306, fig. 4;
- = *Alveolina oblonga* D'ORBIGNY - NUTTALL (1925): 446; pl. 27, fig. 8;
- = *Alveolina oblonga* D'ORBIGNY - HOTTINGER (1960): 141; pl. 9, figs. 4-16.
- Assilina dandotica* DAVIES, 1937
- = *Assilina dandotica* n. sp. - DAVIES in DAVIES & PINFOLD: 28-29; pl. 4, figs. 1-3, 6-8;
- = *Assilina dandotica* DAVIES - SCHAUB (1981): 206-207; pl. 84, figs. 1-16, tab. 18a.
- Assilina daviesi* DE CIZANCOURT, 1938
- = *Assilina daviesi* n. sp. - DE CIZANCOURT: 23; pl. 3, figs. 18, 25-26; tab. 2b (B generation);
- = *Assilina daviesi* var. *nammalensis* n. subsp. - GILL (1953): 82; pl. 13, figs. 1-5 (A generation).
- Assilina lacunata* DE CIZANCOURT, 1946
- = *Assilina umbilicata* n. sp. - DE CIZANCOURT: 23; pl. 3, figs. 19-20, 27 (B generation);
- = *Assilina subumbilicata* n. sp. - DE CIZANCOURT: 24; pl. 3, figs. 21-23 (A generation);
- = *Assilina lacunata* nom. nov. - DE CIZANCOURT (1946): 335;
- Assilina laminosa* GILL, 1953
- = *Assilina laminosa* n. sp. - GILL: 83; pl. 13, figs. 14-17 (B generation);
- = *Assilina sublaminosa* n. sp. - GILL (1953): 83; pl. 13, figs. 18-19 (A generation).
- Assilina leymeriei* (D'ARCHIAC & HAIME, 1853)
- = *Nammulites leymeriei* n. sp. - D'ARCHIAC & HAIME: 153; pl. 11, fig. 9a-c;
- = *Assilina leymeriei* (D'ARCHIAC & HAIME) - SCHAUB (1981): 196-197; fig. 143; pl. 71, figs. 56-74; pl. 72, figs. 1-53, tab. 16d.
- Assilina pustulosa* DONCIEUX, 1926
- = *Assilina pustulosa* n. sp. - DONCIEUX: 52; pl. 5, figs. 36-43; pl. 6, fig. 1 (B generation);
- = *Assilina subpustulosa* n. sp. - DONCIEUX (1926): 53; pl. 5, fig. 20; pl. 6, figs. 2-3 (A generation);
- = *Assilina pustulosa* DONCIEUX - SCHAUB (1981): 207-208; fig. 115; pl. 84, figs. 28-50, 53, 59-96; tab. 18b.
- Assilina spinosa* DAVIES, 1937
- = *Assilina spinosa* n. sp. - DAVIES in DAVIES & PINFOLD: 31-33; pl. 4, figs. 11-12, 16-17 (B generation);
- = *Assilina subspinosa* n. sp. - DAVIES in DAVIES & PINFOLD (1937): 33-34; pl. 4, figs. 19-20, 23-26 (A generation);
- = *Assilina spinosa* DAVIES - SCHAUB (1981): 196; pl. 71, figs. 53-55.
- Dictyoconoides cooki* (CARTER, 1861)
- = *Conulites cooki* n. sp. - CARTER: 83; pl. 15, figs. 7a-g;
- = *Dictyoconoides cooki* (CARTER) - SMOUT (1954): 59-60; pl. 9, figs. 1-5.
- Dictyokathma simplex* SMOUTH, 1954
- = *Dictyokathma simplex* n. sp. - SMOUTH: 64-66; pl. 8, figs. 1-11.
- Discocyclusa dispansa* (SOWERBY, 1840)
- = *Lycophris dispansa* n. sp. - SOWERBY: 327; pl. 24, figs. 16a-b;
- = *Discocyclusa dispansa* (SOWERBY) - NUTTALL (1926): 157; pl. 7, figs. 1-3, 5;
- = *Discocyclusa dispansa* (SOWERBY) - NAGAPPA (1959): 181; pl. 10, figs. 6-8;
- = *Discocyclusa dispansa* (SOWERBY) - SEN GUPTA (1963): 39; pl. 1, figs. 1-9.
- Discocyclusa dorreei* BAYLISS, 1961
- = *Discocyclusa dorreei* n. sp. - BAYLISS: pl. 22, figs. 1-9. Probably the taxonomical status of this species is questionable.
- Discocyclusa ranikotensis* DAVIES, 1927
- = *Discocyclusa ranikotensis* n. sp. - DAVIES: 281-282; pl. 22, figs. 10-12.
- Discocyclusa sowerbyi* NUTTALL, 1926
- = *Discocyclusa sowerbyi* n. sp. - NUTTALL: 149; pl. 8, figs. 1-3;
- = *Discocyclusa sowerbyi* NUTTALL - NAGAPPA (1959): 181; pl. 11, figs. 1-2;
- = *Discocyclusa sowerbyi* NUTTALL - SEN GUPTA (1963): 41; pl. 2, fig. 2, pl. 3, figs. 1-10.
- Discocyclusa undulata* NUTTALL, 1926
- = *Discocyclusa undulata* n. sp. - NUTTALL: 150; pl. 7, figs. 8-9; pl. 8, fig. 5;
- = *Discocyclusa undulata* NUTTALL - NAGAPPA (1959): 181; pl. 10, figs. 9-10.
- Flosculina globosa* (LEYMERIE, 1846)
- = *Alveolina subpyrenaica* var. *globosa* n. subsp. - LEYMERIE: 359; pl. 13, fig. 10a-c;
- = *Flosculina globosa* (LEYMERIE) - NUTTALL (1925): 435; pl. 23, figs. 1-4; pl. 24, figs. 4-6;
- = *Alveolina globosa* LEYMERIE - HOTTINGER (1960): 80; pl. 3, figs. 15-20.
- Kathina maior* SMOUTH, 1954
- = *Kathina major* n. sp. - SMOUTH: 63-64; pl. 6, figs. 1-10.
- Kathina selveri* SMOUTH, 1954
- = *Kathina selveri* n. sp. - SMOUTH: 62-63; pl. 6, figs. 11-13.
- Lockhartia altispra* SMOUTH, 1954
- = *Lockhartia altispra* n. sp. - SMOUTH: 51-52; pl. 4, figs. 4-6.
- Lockhartia conditi* (NUTTALL, 1926)
- = *Dictyoconoides conditi* n. sp. - NUTTALL: 119; pl. 11, figs. 7-8;
- = *Lockhartia conditi* (NUTTALL) - DAVIES (1932): 408; pl. 2, fig. 7; pl. 4, fig. 7;
- = *Lockhartia conditi* (NUTTALL) - DAVIES & PINFOLD (1937): 47-48; pl. 5, fig. 24;
- = *Lockhartia conditi* (NUTTALL) - SMOUTH (1954): 55; pl. 5, figs. 16-19.
- Lockhartia conica* SMOUTH, 1954
- = *Lockhartia conica* n. sp. - SMOUTH: 53; pl. 4, figs. 1-3.



WEISS, W.: Maastrichtian to Eocene foraminiferal assemblages of Pakistan

Plate 9

Plate 9 Rakhi Nala section, Sulaiman Range. Larger foraminifers.

- Fig. 1 *Orbitoides tissoti* SCHLUMBERGER, *Orbitoides* aff. *media* (D'ARCHIAC) and others. - SmR-15/S 1421; about 8 x; Early Maastrichtian.
- Fig. 2 ?*Rotalia* cf. *trochidiformis* (LAMARCK). - SmR-15a/S 1421; about 46 x; Early Maastrichtian.
- Fig. 3 *Orbitoides tissoti* SCHLUMBERGER, *Omphalocyclus macroporus* (LAMARCK) and others. - SmR-15a/S 1421; about 20 x; Early Maastrichtian.
- Fig. 4 Large miliolids, *Omphalocyclus macroporus* (LAMARCK) (left) and others. - SmR-15a/S 1421; about 23 x; Early Maastrichtian.
- Fig. 5 Quadrilocular nucleoconch (= protoconch, deuteroconch, two auxiliary chambers) surrounded by a thick wall and several epiauxiliary chambers of *Orbitoides tissoti* SCHLUMBERGER. - SmR-15a/S 1421; about 46 x; Early Maastrichtian.
- Fig. 6 *Omphalocyclus macroporus* (LAMARCK) showing close affinities to juvenile specimens of *Orbitoides tissoti* SCHLUMBERGER. - SmR-15a/S 1421; about 46 x; Early Maastrichtian.

Lockhartia hamei (DAVIES, 1927)

- = *Dictyoconoides hamei* n. sp. - DAVIES: 280-281; pl. 21, figs. 13-15;
- = *Lockhartia hamei* (DAVIES) - DAVIES (1932): 407; pl. 2, figs. 4-6;
- = *Lockhartia hamei* (DAVIES) - DAVIES & PINFOLD (1937): 45-46; pl. 7, figs. 9-13, 15;
- = *Lockhartia hamei* (DAVIES) - SMOOTH (1954): 49-50; pl. 2, figs. 1-14.

Lockhartia hunti OVEY, 1947

- = *Lockhartia hunti* n. sp. - OVEY: 573; pl. 10, figs. 1-6; pl. 11, fig. 1;
- = *Lockhartia hunti* OVEY - SMOOTH (1954): 54; pl. 4, fig. 7.

Lockhartia hunti pustulosa SMOOTH, 1954

- = *Lockhartia hunti* var. *pustulosa* n. subsp. - SMOOTH: 54-55; pl. 4, figs. 8-10.

Lockhartia tipperi (DAVIES, 1926)

- = *Conulites tipperi* n. sp. - DAVIES: 247-248; pl. 18, fig. 8;
- = *Lockhartia tipperi* (DAVIES) - DAVIES (1932): 407;
- = *Lockhartia tipperi* (DAVIES) - DAVIES in DAVIES & PINFOLD (1937): 48-49; pl. 6, figs. 14-16; pl. 7, fig. 17;
- = *Lockhartia tipperi* (DAVIES) - SMOOTH (1954): 55; pl. 4, figs. 11-13.

Miscellanea miscella (D'ARCHIAC & HAIME, 1854)

- = *Nummulites miscella* n. sp. - D'ARCHIAC & HAIME: 345; pl. 35, figs. 4a-c;
- = *Siderolites stampeii* n. sp. - DAVIES (1927): 278; pl. 21, figs. 1-8;
- = *Miscellanea miscella* (D'ARCHIAC & HAIME) - PFENDER (1935): 231-235, figs. 1-4; pl. 11, figs. 6-7; pl. 13, figs. 2-4;
- = *Miscellanea miscella* (D'ARCHIAC & HAIME) - DAVIES & PINFOLD (1937): 43-45; pl. 6, figs. 1-3, 5, 7-8.

Nummulites atacicus LEYMERIE, 1846

- = *Nummulites atacica* n. sp. - LEYMERIE (1846): 358; pl. 13, figs. 13a-e;
- = *Nummulites atacicus* LEYMERIE - H. DOUVILLÉ (1919): 38, fig. 2; pl. 3, figs. 3-6 (B generation);
- = *Nummulites subatacicus* n. sp. - H. DOUVILLÉ (1919): 41, fig. 3; pl. 3, figs. 7-8 (A generation);
- = *Nummulites atacicus* LEYMERIE - SCHAUB (1981): 119-120; pl. 25, figs. 1-51; tab. 14i.

Nummulites crasseornata (HENRICI, 1934)

- = *Camerina crasseornata* n. sp. - HENRICI: 32, 52; pl. 2, figs. 2-5, 12; text-fig. 15.

Nummulites fossulata DE CIZANCOURT, 1946

- = *Nummulites fossulata* n. sp. - DE CIZANCOURT: 644, 648, 654; pl. 10, figs. 6-8; text-fig. 1.

Nummulites globulus LEYMERIE, 1846

- = *Nummulites globulus* n. sp. - LEYMERIE (1846): 359; pl. 13, figs. 14a-d;
- = *Nummulites globulus* LEYMERIE - SCHAUB (1981): 137-138; pl. 40, figs. 1-80; tab. 1f.

Nummulites vanderstoki RUTTEN & VERMUNT, 1932

- = *Nummulites vanderstoki* n. sp. - RUTTEN & VERMUNT: 240; pl. 1, fig. 8a-b; pl. 2, figs. 6, 12.

Operculina aspensis COLOM, 1954

- = *Operculina aspensis* n. sp. - COLOM: 183; pl. 10, figs. 6-8.
- = *Operculina aspensis* COLOM - HOTTINGER (1977): 95; fig. 37o-v.

Operculina canalifera D'ARCHIAC & HAIME, 1853

- = *Operculina canalifera* D'ARCHIAC & HAIME - NUTTALL (1926): 117; pl. 11, figs. 1-2.

Operculina patalaensis DAVIES, 1937

- = *Operculina patalaensis* n. sp. - DAVIES in DAVIES & PINFOLD (1937): 38-39; pl. 5, figs. 6, 17, 19, 26 (B generation).
- = *Operculina jiwani* n. sp. - DAVIES in DAVIES & PINFOLD (1937): 39-40; pl. 5, figs. 11-14, 16 (A generation).

Operculina salsa DAVIES, 1937

- = *Operculina salsa* n. sp. - DAVIES in DAVIES & PINFOLD (1937): 37; pl. 5, figs. 1, 3, 7, 10, 15 (B generation);
- = *Operculina subsalsa* n. sp. - DAVIES in DAVIES & PINFOLD (1937): 37; pl. 5, figs. 2, 4, 8-9 (A generation).

Orbitolites complanatus LAMARCK, 1801

- = *Orbitolites complanatus* n. sp. - LAMARCK: 376;
- = *Orbitolites complanata* LAMARCK - NUTTALL (1925): 447;
- = *Orbitolites complanata* LAMARCK - DAVIES (1937): 66.

Ranikothalia sp.

Under this genus specimens were united which were either close to *Operculina sindensis* DAVIES, 1927, and *Operculina bermudezi* PALMER, 1934, which are regarded as *Ranikothalia sindensis* (DAVIES), or close to *Nummulites nuttalli* DAVIES, 1927, which is regarded as *Ranikothalia nuttalli* (DAVIES) (compare HOTTINGER 1977: 50-52, BUTT 1987, 1991).

Rotalia trochidiformis (LAMARCK, 1804)

- = *Rotalites trochidiformis* n. sp. - LAMARCK: 183-185;
- = *Dictyoconoides newboldi* (D'ARCHIAC & HAIME, 1854) - DAVIES (1927): 279; pl. 22, figs. 1-4;
- = *Rotalia trochidiformis* (LAMARCK) - SMOOTH (1954): 43-45; pl. 1, figs. 1-6.